

**CAUTION** KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



# Chlorsulfuron

## 750 WG

## Herbicide

**ACTIVE CONSTITUENT: 750g/kg CHLORSULFURON**

**GROUP B HERBICIDE**

A selective herbicide for the control of annual (wimmera) ryegrass and certain broadleaf weeds in wheat, barley, oats and cereal rye and triticale

**IMPORTANT: READ THE ATTACHED LEAFLET  
BEFORE OPENING OR USING THIS PRODUCT**

CONTENTS

**1kg**

### STORAGE AND DISPOSAL KEEP OUT OF REACH OF CHILDREN

Store in the closed, original container in a well ventilated area, as cool as possible. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty containers for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty containers 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory Government Regulations. DO NOT burn empty containers or product.

### SAFETY DIRECTIONS

Avoid contact with eyes and skin. DO NOT inhale spray mist. Wash hands after use.

### FIRST AID

If poisoning occurs contact a Doctor or phone the Australian Poisons Information Centre 13 11 26.

### MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS) that can be obtained from the supplier.



**Wellfarm Pty Ltd**  
ACN 158 518 551  
ABN 32 158 518 551

22 Calypso Crescent  
Point Cook Vic 3030  
[www.wellfarm.com.au](http://www.wellfarm.com.au)

### CONDITIONS OF SALE

The use of this product is beyond the control of the manufacturer. No warranty expressed or implied is given by Wellfarm Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Wellfarm Pty Ltd accepts NO responsibility for any consequences whatsoever resulting from the use of this product.

### MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY

Wellfarm Pty Ltd has no control over storage, handling and manner of use of this product. Where the material is not stored, handled or used correctly and in accordance with the directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Wellfarm Pty Ltd accepts NO liability for any losses or damages arising from incorrectly storage, handling or use.

**APVMA Approval No 68337/58155**

**IN AN EMERGENCY  
DIAL 000 POLICE  
OR FIRE BRIGADE**

Batch No

DOM



# CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

## Wellfarm Chlorsulfuron 750 WG Herbicide

ACTIVE CONSTITUENT: 750g/kg CHLORSULFURON

GROUP	<b>B</b>	HERBICIDE
-------	----------	-----------

A selective herbicide for the control of annual (wimmera) ryegrass and certain broadleaf weeds in wheat, barley, oats and cereal rye and triticale

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE  
OPENING OR USING THIS PRODUCT**

**CONTENTS: 1kg  
(500g)**

APVMA Approval No 68337/58155

Wellfarm Pty Ltd  
ACN 158 518 551  
22 Calypso Crescent  
Point Cook Vic 3030  
[www.wellfarm.com.au](http://www.wellfarm.com.au)



**DIRECTIONS FOR USE****RESTRAINTS**

DO NOT spray emerged crops if rain is expected within 4 hours.

After mixing in the tank, spray within 48 hours if Chlorsulfuron is used by itself, or within 24 hours if mixed with another product.

DO NOT apply to plants suffering stress.

**METHOD OF USE – PRE-SOWING INCORPORATED BY SOWING****ANNUAL RYEGRASS**

ANNUAL RYEGRASS						
Crop Situation	Weeds Controlled	State(s)	Rate g/ha			Critical Comments
			Soil Type			
			Light to Medium Soils		Heavy Soils	
			Soil pH			
			Less than 7	7.0 – 8.5	8.5 or less	
Wheat and Triticale only	Annual (Wimmera) Ryegrass <i>Lolium rigidum</i>	NSW, ACT, VIC, SA, WA only	20	15 or 20*	20	* Use the higher rate when paddock history suggests a high weed population can be expected.  NOTE: Refer to General Instructions for optimum application timing and conditions.

Crop Situation	Weeds Controlled	State(s)	Rate g/ha	Critical Comments
Wheat and Triticale only	African Turnip Weed Sisymbrium thellungii	NSW, ACT and QLD only	20	If possible, spray and incorporate into the soil in one operation, If this is not possible, incorporation should take place within 4 hours of spraying. Delay may cause inferior weed control.
	Amsinckia/Yellow Burrweed Amsinckia spp.	NSW, ACT, VIC, SA, WA only	15	
	Annual; Phalaris Phalaris paradoxa Phalaris minor	NSW, ACT only	20 plus 830 mL trifluralin (480g/L)	
	Barley Grass Hordeum leporinum	NSW, ACT and TAS only		
	Silver grass Vulpia spp.	TAS only		
	Ball Mustard Neslia puniculata	SA only	15	
	Black Bindweed/ Climbing Buckwheat Fallopia convolvulus	QLD only	20	Apply to dry soil before the sowing rain. Mechanical incorporation before the sowing rains is NOT necessary.
	Brome grass Bromus spp.	NSW, ACT, VIC, SA,	20	Gives suppression only if populations are 20 plants/m <sup>2</sup> or less.

Crop Situation	Weeds Controlled	State(s)	Rate g/ha	Critical Comments
	(suppression only)	WA, TAS only		
	Cape Tulip Homeria spp.	WA only		On acid soils pH 5.5 or less, this product will give a shorter period of control in wet years.
	Capeweed Arctotheca calendula	NSW, ACT, VIC, SA, WA, TAS only		
	Charlock Sinapis arvensis	VIC, SA, TAS only	15	
	Common Iceplant Mesembryanthemum crystallinum	SA only		
	Corn Gromwell, Sheepweed, White Ironweed Buglossoides arvensis	QLD, NSW, ACT, VIC, SA, WA only	20	
	Deadnettle Lamium amplexicaule	All states	15 or 20	Use the higher rate when paddock history suggests a high weed population can be expected.
	Docks Rumex spp.	NSW, ACT, VIC, SA, WA, TAS only	20	
	Fat-Hen Chenopodium album	NSW, ACT TAS only		
	Fumitory Fumaria spp.	NSW, ACT, VIC, SA, WA, TAS only	15 or 20	Use the higher rate when paddock history suggests a high weed population can be expected.
	Guildford Grass/Onion grass Romulea rosea	WA only	15	
	Indian Hedge Mustard Sisymbrium oriental	All states		
	King Island Melilot Melilotus indicus	VIC, SA only		
	Lincoln Weed Diplotaxis tenuifolia	SA only		
	Loosestrife Lysimachia spp	VIC only		

Crop Situation	Weeds Controlled	State(s)	Rate g/ha	Critical Comments
	Mintweed <i>Salvia reflexa</i>	QLD, NSW, ACT only	20	Apply to dry soil before the sowing rain. Mechanical incorporation before the sowing rains is NOT necessary.
	Mouse-Ear Chickweed <i>Cerastium</i> spp.	NSW, ACT, VIC, SA WA, TAS only	15	
	New Zealand Spinach <i>Tetragonia tetragonoides</i>	QLD only	20	
	Paradoxa Grass <i>Phalaris paradoxa</i>	Nth NSW (soil pH > 7.5) and QLD only		
	Paterson's Curse/Salvation Jane <i>Echium plantagineum</i>	NSW, ACT, VIC, SA, WA, TAS only	15	
	Pimpernels <i>Anagallis arvensis</i>	NSW, ACT, VIC, SA, TAS only		
	Prickly Lettuce/Whip Thistle <i>Lactuca serriola</i>	VIC, SA only	20	
	Rough Poppy <i>Papaver hybridum</i>	NSW, ACT, SA, WA, TAS only	15 or 20	Use the higher rate when paddock history suggests a high weed population can be expected.
	Saffron Thistle (suppression only) <i>Carthamus lanatus</i>	QLD, NSW, ACT, VIC, SA, TAS only	20	
	Saltbush <i>Atriplex muelleri</i>	QLD, NSW, ACT only		
	Shepherd's Purse <i>Capsella bursa-pastoris</i>	NSW, ACT, VIC, SA, WA, TAS only	15 or 20	Use the higher rate when paddock history suggests a high weed population can be expected.
	Slender Celery <i>Apium leptophyllum</i>	QLD, NSW, ACT only	20	
	Slender Thistle <i>Carduus tenuiflorus</i>	TAS only		
	Soursob <i>Oxalis pes-caprae</i>	NSW, ACT, VIC, SA only	15	Apply only to soils of pH 7.5 or above. Apply after majority of soursobs have emerged and leave soil undisturbed for 1–4 weeks prior to cultivating or sowing. The most effective and reliable control is achieved with early post-emergence

Crop Situation	Weeds Controlled	State(s)	Rate g/ha	Critical Comments
				applications (EPE) after crop and weed emergence.
	Spear Thistle <i>Cirsium vulgare</i>	TAS only	20	
	Stemless Thistle <i>Onopordum acaulon</i>	SA only	15 or 20	Use the higher rate when paddock history suggests a high weed population can be expected.
	Storksbill/Wild Geranium <i>Erodium</i> spp.	VIC, SA, WA, TAS only	15	
	Three cornered Jack(s) /Doublegee/Spiny Emex <i>Emex australis</i>	NSW, ACT, VIC, SA, WA only	20	
	Tree Hogweed <i>Polygonum patulum</i>	VIC, SA only		
	Turnip Weed <i>Rapistrum rugosum</i>	QLD and SA only	15	
	Wireweed/Hogweed <i>Polygonum aviculare</i>	All states	15 or 20	Use the higher rate when paddock history suggests a high weed population can be expected.
	Wild Turnip <i>Brassica tournefortii</i>	NSW, ACT, VIC, SA, WA, TAS only	15	

**METHOD OF USE – POST CROP AND WEED EMERGENCE**  
**ANNUAL RYEGRASS**

ANNUAL RYEGRASS						
Crop Situation	Weeds Controlled	State(s)	Rate g/ha			Critical Comments
			Soil Type			
			Light to Medium Soils		Heavy Soils	
			Soil pH			
			Less Than 7	7.0 – 8.5	8.5 or Less	
Wheat, Barley, Oats, Cereal Rye and Triticale only	Annual (Wimmera) Ryegrass <i>Lolium rigidum</i>	NSW, ACT, VIC, SA, WA only	20 or 25*	15 or 20*	20 or 25*	<p>* Use the higher rate when paddock history suggests a high weed population can be expected.</p> <p>Application of this product to Annual Ryegrass 2 leaf or greater with water volumes less than 50L/ha may result in reduced efficacy.</p>

Crop Situation	Weeds Controlled	State(s)	Rate g/ha	Critical Comments
Wheat, Barley, Oats, Cereal Rye and Triticale only	African Turnip Weed Sisymbrium thellungii	NSW, ACT and QLD only	20	Apply at cotyledon to 4 leaf stage.
	Amsinckia/Yellow Burrweed Amsinckia spp.	NSW, ACT, VIC, SA, WA only	15	
	Ball Mustard Neslia puniculata	SA only	25	
	Bifora/Carrot Weed Cotula australis			
	Black Bindweed/ Climbing Buckwheat Fallopia convolvulus	QLD, NSW, ACT only	20	Apply at cotyledon to 2 leaf stage of weed.
	Cape Tulip Homeria spp.	WA only	15	
	Charlock Sinapis arvensis	NSW, ACT, VIC, SA, TAS only		
	Corn Gromwell, Sheepweed, White Ironweed Buglossoides arvensis	NSW, ACT, VIC, SA, WA only	20	Apply at cotyledon to 2 leaf stage of weed. If applied at a later stage only suppression will occur.
	Deadnettle Lamium amplexicaule	QLD, NSW, ACT, VIC, SA, TAS only	15 or 20	Use the higher rate under heavy weed pressure.

Crop Situation	Weeds Controlled	State(s)	Rate g/ha	Critical Comments
	Docks Rumex spp.	VIC, SA, WA, TAS only	15	Apply at cotyledon to 2 leaf stage.
	Fat-Hen Chenopodium album	NSW, ACT TAS only	20	
	Fumitory, Denseflower Fumaria spp.	NSW, ACT, VIC, SA, WA, TAS only		
	Guildford Grass/Onion grass Romulea rosea	WA only	15	
	Hoary Cress Cardaria draba	VIC, SA, TAS only	20	Apply when plants are fully emerged.
	Lincoln Weed Diplotaxis tenuifolia	SA only		
	Matricaria Matricaria matricarioides	WA, TAS only		
	Mintweed Salvia reflexa	QLD, NSW, ACT only		Apply at cotyledon to 4 leaf stage.
	Mouse-Ear Chickweed Cerastium spp.	NSW, ACT, VIC, SA, WA, TAS only	15	
	Mustards Sisymbrium spp.	All states		
	New Zealand Spinach Tetragonia tetragonoides	QLD only	20	
	Paterson's Curse/Salvation Jane Echium plantagineum	NSW, ACT, VIC, SA, WA, TAS only	15	
	Pimpernels Anagallis arvensis	NSW, ACT, VIC, SA, TAS only		
	Prickly Lettuce/Whip Thistle Lactuca serriola	VIC, SA only	20	
	Rough Poppy Papaver hybridum	NSW, ACT, SA, WA, TAS only		



Crop Situation	Weeds Controlled	State(s)	Rate g/ha	Critical Comments
	Saltbush <i>Atriplex Muelleri</i>	QLD, NSW, ACT only		Apply at cotyledon to 4 leaf stage.
	Shepherd's Purse <i>Capsella bursa-pastoris</i>	NSW, ACT, VIC, SA, WA, TAS only		
	Slender Celery <i>Apium leptophyllum</i>	QLD, NSW, ACT only		Apply at cotyledon to 4 leaf stage.
	Soursob <i>Oxalis pes-caprae</i>	NSW, ACT, VIC, SA only		Apply when the majority of soursobs have emerged.
	Spear Thistle <i>Cirsium vulgare</i>	TAS only		
	Stagger weed <i>Stachys arvensis</i>	QLD, NSW, ACT, WA, TAS only		
	Stemless Thistle <i>Onopordum acaulon</i>	VIC only	25	
	Storksbill/Wild Geranium <i>Erodium spp.</i>	VIC, SA, WA, TAS only	15	
	Tree Hogweed <i>Polygonum patulum</i>	VIC only	20	
	Turnip Weed <i>Rapistrum rugosum</i>	All states	15	
	Wild Radish <i>Raphanus raphanistrum</i>	All states	15 or 20	Use the higher rate under heavy weed pressure. A follow-up spray with a suitable herbicide may be necessary to control subsequent germinations.
	Wild Turnip <i>Brassica tournefortii</i>	NSW, ACT, VIC, SA, WA, TAS only	15	
	Wireweed/Hogweed <i>Polygonum aviculare</i>	All states	20	

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL  
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIODS: NOT REQUIRED WHEN USED AS DIRECTED.**

## GENERAL INSTRUCTIONS

This product is a selective herbicide designed to control certain weeds in wheat, triticale, barley, oats and cereal rye.

This product is suitable as a pre-sowing treatment for wheat and triticale, and as a post-sowing treatment for wheat, triticale, barley, oats and cereal rye. When used on emerged weeds, the product is absorbed by foliage and green stems and moves into the root system.

Prior to using this product, careful consideration should be given to soil pH. As soil pH increases, rate of breakdown decreases.

This product should NOT be used on soils with a pH of 8.6 or higher as soil residual activity could adversely affect following crops and crop rotation intervals may be extended beyond normal intervals.

Crops other than wheat, barley, oats, triticale and cereal rye can be extremely sensitive to low concentrations of this product in the soil. See Crop Rotation Recommendations.

Best weed control is obtained when rainfall or sprinkler irrigation wets the soil to a depth of 5 to 7.5 cm within 4 weeks of application.

### Pre-Sowing Incorporated by Sowing:

WA only – Avoid applying to dry sandy soils as rapid leaching may occur with early season rains.

SA – Before using rates greater than 15g/ha on light to medium soils pH 7 to 8.5, seek further advice.

Conventionally Sown Crops – On soils less than pH 7, apply a spray just before sowing or in conjunction with the sowing operation. On soils of pH 7 or greater it is NOT critical to time the spray just before sowing. Spray onto non-ridged surface free of large clods. Use low profile 10cm combine points for sowing. Sow at speeds of 10kph or greater. Use light covering harrows at sowing. If applied to dry soil and sowing is to be delayed, incorporate immediately after spraying to prevent loss by wind erosion.

**Direct Drilled Crops** – Apply tank mixed with either paraquat/diquat mixture of glyphosate in accordance with manufacturer's label recommendations.

### Post Crop and Weed Emergence:

Where treatment is delayed or where weeds are NOT actively growing due to adverse conditions results may be slow to appear and weeds may be only stunted or suppressed.

**Wheat, Triticale, and Cereal Rye** – Apply after crop emergence and when weeds are small and actively growing (Annual Ryegrass no more than 3 leaves), broadleaved weeds no more than 5cm in height or diameter (for Black Bindweed refer to Specific Recommendations).

**Barley and Oats** – Apply between the 2 leaf stage of the crop (3 leaf stage in SA) and early tillering, when weeds are small and actively growing. (Annual Ryegrass no more than 3 leaves), Broadleaved weeds NO more than 5cm in height or diameter (for Black Bindweed refer to Specific Recommendations).

## RESISTANT WEEDS WARNING

<b>GROUP</b>	<b>B</b>	<b>HERBICIDE</b>
--------------	----------	------------------

Wellfarm Chlorsulfuron 750 WG Herbicide is a member of the sulfonyleurea group of herbicides.

Wellfarm Chlorsulfuron 750 WG Herbicide has the inhibitor of the enzyme acetolactate synthase (ALS) mode of action. For weed resistance management, Wellfarm Chlorsulfuron 750 WG Herbicide is a Group B herbicide.

Some naturally occurring weed biotypes resistant to Wellfarm Chlorsulfuron 750 WG Herbicide and other Group B herbicides (Annual Ryegrass and some broadleaf weeds) may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if Wellfarm Chlorsulfuron 750 WG Herbicide and other Group B herbicides (Annual Ryegrass and some broadleaf weeds) are used repeatedly. These resistant weeds will NOT be controlled by Wellfarm Chlorsulfuron 750 WG Herbicide or any other Group B herbicides (Annual Ryegrass and some broadleaf weeds).

Annual Ryegrass biotypes resistant to diclofop-methyl and other "grass specific" herbicides are often also resistant to Wellfarm Chlorsulfuron 750 WG Herbicide. Before using Wellfarm Chlorsulfuron 750 WG Herbicide on a population resistant to "grass specific" herbicides, have a resistance test conducted to ensure that it is still susceptible to Wellfarm Chlorsulfuron 750 WG Herbicide.

Since the occurrence of resistant weeds is difficult to detect prior to use, Wellfarm Pty Ltd accepts NO liability for any losses that may result from the failure of Wellfarm Chlorsulfuron 750 WG Herbicide to control resistant weeds.

To prevent, or at least minimise the risk of resistant weeds occurring, use Wellfarm Chlorsulfuron 750 WG Herbicide in tank mixes (if appropriate) and/or rotations with herbicides having different modes of action effective on the same weed species.

Large numbers of healthy surviving weeds can be an indication that resistance is developing. Efforts should be taken to prevent seed set of these survivors.

DO NOT make more than one application of an ALS inhibitor herbicide to a crop, either pre-sowing incorporated by sowing or post crop and weed emergence.

If the user suspects that an ALS inhibitor-resistant weed is present, Wellfarm Chlorsulfuron 750 WG Herbicide or other ALS inhibitor herbicides should NOT be used.

Wellfarm Chlorsulfuron 750 WG Herbicide may be subject to specific resistance management strategies. For further information consult your local farm chemical supplier, consultant, local Department of Agriculture or Primary Industries.

### **GRAZING ADVICE**

Avoid grazing treated areas within 24 hours of application to optimise weed control.

A nil withholding period is applicable for grazing Wellfarm Chlorsulfuron 750 WG Herbicide treated areas (when used as directed on the label).

### **CROP SAFETY**

DO NOT use this product for:

- crops other than cereals.
- cereals irrigated by furrows or flooding.
- winter cereals undersown with legume pasture crops.
- weed control where crops are under stress. Damage can occur where crops are stressed due to conditions such as excessive soil alkalinity or acidity, poor nutrient status, disease, nematode or insect infestation, adverse weather conditions, drought or waterlogging. If crops become stressed after spraying, they may turn yellow or become retarded, but usually they will recover with no reduction in yield.

### **WHEAT**

DO NOT use this product for:

- wheat varieties Cranbrook, or Miling.
- the wheat variety Vulcan if on acid soils and under stress conditions caused by waterlogging, frost, aluminium or manganese toxicity; reduced yields may result.
- pre-sowing treatment of weeds in wheat varieties Avocet and Durati (okay for post-emergent use).
- pre-sowing treatment of weeds in wheat variety Banks if soil pH is 5.5 or less (okay for post-emergent use).

### **BARLEY AND OATS**

DO NOT use this product for:

- application before the crop has reached the 2 leaf stage (3 leaf stage in SA).
- Stirling barley.
- Barley under waterlogged conditions (yield may be reduced).

The application of other sulfonylurea herbicides following this product is NOT recommended.

### **CROP ROTATION RECOMMENDATIONS**

Land previously treated with this product should NOT be rotated to crops other than those listed in the following tables.

Tolerance of other crops (grown through to maturity) should be determined on a small scale before sowing into larger areas.

The treated areas may be re-planted to any of the specified crops after the interval indicated in the following tables:

### **NB – THE TABLE BELOW APPLIES TO ALL STATES**

MINIMUM RECROPPING INTERVAL (Months After Application)						
	0	3	6	9	12	18
<b>Soil pH*</b>	Triticale Wheat	Cereal Rye	Oats	Barley	Subterranean Clover ** Faba Beans Field Pea	Maize Sorghum Soybeans Sunflower
6.5 or less						

					Linseed Lucerne Lupins Medics ** Rapeseed Safflower	
--	--	--	--	--	--	--

**NB – THE TABLES BELOW APPLY TO QLD, SA, WA & TAS ONLY**

AND THE TABLES BELOW APPLY TO QED, CA, WA & TAS ONLY

MINIMUM RECROPPING INTERVAL (Months After Application)						
Rainfall Requirement	0	3	9	15	18	22
	Minimum 700mm					
Soil pH*	Triticale Wheat	Cereal Rye	Barley Oats	Japanese Millet Maize Panicum Millet Sorghum Sunflower White French Millet	Cotton Soybeans	Faba Beans Field Pea Linseed Medics ** Rapeseed Safflower Subterranean Clover **
6.6 to 7.5						
MINIMUM RECROPPING INTERVAL (Months After Application)						
Rainfall Requirement	0	15	18	24 months or longer		
	Minimum 700mm					
Soil pH*	Triticale Wheat	Japanese Millet Maize Panicum Millet Sorghum Sunflower White French Millet	Barley Oats Cereal Rye	Rotate to crops other than Cereals (such as listed above) only if field test strip of planned rotational crop has been successfully grown to maturity in the previous season.		
7.6 to 8.5						
Soil pH*	This product is NOT recommended for use on soils of pH 8.6 and above.					
8.6 and above						

**NB – THE TABLES BELOW APPLY TO NSW, ACT & VIC ONLY**

ND – THE TABLES BELOW APPLY TO NOW, AOT & VIC ONLY

MINIMUM RECROPPING INTERVAL (Months After Application)					
	0	3	9	22	29
<b>Soil pH*</b>  6.6 to 7.5	Triticale Wheat	Cereal Rye	Barley Oats	Faba Beans Field Pea Linseed Lucerne Lupins Medics ** Subterranean Clover **	Maize Sorghum Soybeans Sunflower
MINIMUM RECROPPING INTERVAL (Months After Application)					
	0	18	24 months or longer		
<b>Soil pH*</b>  7.6 to 8.5			Rotate to crops other than Cereals (such as listed above) only if field test strip of planned rotational crop has been successfully grown to maturity in the previous season.		
<b>Soil pH*</b>  8.6 and above	This product is NOT recommended for use on soils of pH 8.6 and above.				

\* Soil pH is determined by laboratory analysis using the 1:5 soil:water suspension method.

\*\* Include natural regeneration of Subterranean clover and medics.

- Land previously treated with this product should NOT be rotated to crops other than those listed in the above table.

- **Tolerance of other crops (grown through to maturity) should be determined on a small scale before sowing into larger areas.**

## **SPRAY PREPARATION**

Wellfarm Chlorsulfuron 750 WG Herbicide is a water dispersible granule.

1. Fill tank partially with water and engage full agitation.
2. Add the required amount (N.B. The measuring flask provided is graduated in grams of Wellfarm Chlorsulfuron 750 WG Herbicide only. DO NOT use for measuring of other materials).
3. Top up with water to the required volume.
4. Companion Products: If applying this product with another product ensure this product is completely dissolved before adding the companion product.
5. Wellfarm Chlorsulfuron 750 WG Herbicide must be kept in suspension at all times by continuous agitation. Where prepared spray mixes have been allowed to stand, thoroughly re-agitate before using.

## **USE OF SURFACTANT/WETTING AGENT**

For post emergent application always add a non-ionic surfactant at 100mL per 100L of final spray volume (0.1% volume/volume).

The use of spraying oils is NOT recommended.

**NOTE:** DO NOT add surfactant/wetting agent when product is tanked with another product that already has a surfactant/wetting agent in the formulation.

## **SPRAYING**

### **GROUND APPLICATION**

Use a boom spray properly calibrated to a constant speed and rate of delivery to ensure thorough coverage and a uniform spray pattern. Avoid overlapping and shut off spray booms while starting, turning, slowing or stopping as injury to the crop may result. Apply a minimum of 30L of spray mix per hectare.

### **AERIAL APPLICATION**

Apply at minimum of 20L/ha water. Avoid spraying in still conditions or in winds likely to cause drift onto adjacent sensitive crops. Avoid spraying where drift can go onto areas likely to be sown to sensitive crops – see Crop Rotation Recommendations. Turn off spray boom whilst passing over creeks and dams.

## **SPRAYER CLEAN-UP**

It is essential that the sprayer be properly cleaned after using this product to prevent injury to crops other than wheat, triticale, barley, oats or cereal rye. All traces of Wellfarm Chlorsulfuron 750 WG Herbicide should be removed from equipment using the following procedure:

1. Drain tank, then flush tank, boom and hoses with clean water for at least 10 minutes.
2. Fill tank with clean water then add 300 mL of household chlorine bleach (4% chlorine) per 100L of water. Flush through boom and hoses, then allow to sit for 15 minutes with agitation engaged, then drain.
3. Repeat steps 2.
4. Nozzles and screen should be removed and cleaned separately. To remove traces of chlorine bleach, rinse the tank thoroughly with clean water and flush through hoses and boom.

## **CAUTION**

DO NOT use chlorine bleach with ammonia. All traces of liquid fertiliser contacting ammonia, ammonium nitrate or ammonium sulphate must be rinsed with water from mixing and application equipment before adding chlorine bleach solution. Failure to do so will release a gas with a musty chlorine odour, which can cause eye, nose, throat and lung irritation. DO NOT clean equipment in an enclosed area.

## **COMPATIBILITY**

**Wellfarm Chlorsulfuron 750 WG Herbicide Is Compatible With Glyphosate And Paraquat. The Product Does NOT Control Wild Oats, However It Is Compatible With Wild Oat Herbicides: Tri-Allate, Flamprop-M-Methyl And Fenoxaprop-P-Ethyl. It Is Also Compatible With Trifluralin And; Broadleaf Herbicides:**

Diflufenican/MCPA, Diflufenican/Bromoxynil, 2,4-D Amine, 2,4-D Ester, Bromoxynil, MCPA, and Bromoxynil/MCPA Mixtures.

**Insecticides:**

Chlorpyrifos, Omethoate, Dimethoate, Deltamethrin and Fenvalerate.

#### **PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

DO NOT apply or drain or flush equipment on or near desirable trees or other plants or on areas where their roots may extend or in locations where the chemical may be washed or moved into contact with their roots.

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pasture.

#### **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate dams, streams, drains, rivers or waterways with the chemicals or used containers.

#### **STORAGE AND DISPOSAL**

##### **KEEP OUT OF REACH OF CHILDREN**

Store in the closed, original container in a well ventilated area, as cool as possible. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty containers for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty containers 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory Government Regulations. DO NOT burn empty containers or product.

#### **SAFETY DIRECTIONS**

Avoid contact with eyes and skin. DO NOT inhale spray mist. Wash hands after use.

#### **FIRST AID**

If poisoning occurs contact a Doctor or phone the Australian Poisons Information Centre 13 11 26.

#### **MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet (MSDS) that can be obtained from the supplier.

#### **CONDITIONS OF SALE**

The use of this product is beyond the control of the manufacturer. No warranty expressed or implied is given by Wellfarm Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Wellfarm Pty Ltd accepts NO responsibility for any consequences whatsoever resulting from the use of this product.

#### **MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY**

Wellfarm Pty Ltd has no control over storage, handling and manner of use of this product. Where the material is not stored, handled or used correctly and in accordance with the directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Wellfarm Pty Ltd accepts NO liability for any losses or damages arising from incorrectly storage, handling or use.

**IN AN EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE**

APVMA Approval No 68337/58155

Wellfarm Pty Ltd  
ACN 158 518 551  
22 Calypso Crescent  
Point Cook Vic 3030  
[www.wellfarm.com.au](http://www.wellfarm.com.au)