

CAUTION KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Fluroxypyr 200 EC

Herbicide

**ACTIVE CONSTITUENT: 200g/L FLUROXYPYR
PRESENT AS THE METHYL HEPTYL ESTER
SOLVENT: 586g/L HYDROCARBON LIQUID**

GROUP

I

HERBICIDE

For the control of a wide range of broadleaf weeds in fallow, lucerne, maize, millets, pastures, sorghum, sugar cane, sweet corn, winter cereals. Also for the control of woody weeds in agricultural non-crop areas, commercial and industrial areas, pastures and rights-of-way as specified in the Directions For Use Table

**IMPORTANT: READ THE ATTACHED LEAFLET
BEFORE OPENING OR USING THIS PRODUCT**

CONTENTS

20L

STORAGE AND DISPOSAL KEEP OUT OF REACH OF CHILDREN

Storage For All Containers

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight.

Disposal Recycled Containers

This container can be recycled if it is clean, dry, free of visible residues and has the drum MUSTER logo visible. Triple or preferably pressure rinse containers before disposal. Dispose of rinse by adding to the spray tank. DO NOT dispose of undiluted chemicals on site. Wash outside of the container and cap. Stored cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at a drum MUSTER collection or similar container management site. The cap should NOT be replaced but taken separately.

Disposal Non-Recycled Containers

Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty containers for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty containers 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory Government Regulations. DO NOT burn empty containers or product.

Disposal Refillable Containers

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SMALL SPILL MANAGEMENT

Wear protective equipment (see SAFETY DIRECTIONS). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal (see STORAGE AND DISPOSAL section). If necessary wash the spill area with an alkali detergent and water and absorb the wash liquid for disposal.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. Sensitive workers should use protective clothing when opening the container and preparing the spray and using the prepared spray.



Wellfarm Pty Ltd
ACN 158 518 551
ABN 32 158 518 551

22 Calypso Crescent
Point Cook Vic 3030
www.wellfarm.com.au

Wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and face shield or goggles. After each day's use, wash gloves, face shield or goggles and contaminated clothing. Wash hands after use.

FIRST AID

If poisoning occurs contact a Doctor or phone the Australian Poisons Information Centre 13 11 26.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS) that can be obtained from the supplier.

CONDITIONS OF SALE

The use of this product is beyond the control of the manufacturer. No warranty expressed or implied is given by Wellfarm Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Wellfarm Pty Ltd accepts NO responsibility for any consequences whatsoever resulting from the use of this product.

MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY

Wellfarm Pty Ltd has no control over storage, handling and manner of use of this product. Where the material is not stored, handled or used correctly and in accordance with the directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Wellfarm Pty Ltd accepts NO liability for any losses or damages arising from incorrectly storage, handling or use.

APVMA Approval No 70149/62623

**IN AN EMERGENCY
DIAL 000 POLICE
OR FIRE BRIGADE**



Batch No

DOM

CAUTION

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

Wellfarm Fluroxypyr 200 EC Herbicide

**ACTIVE CONSTITUENT: 200g/L FLUROXYPYR
PRESENT AS THE METHYL HEPTYL ESTER
SOLVENT: 586g/L HYDROCARBON LIQUID**

GROUP	I	HERBICIDE
--------------	----------	------------------

For the control of a wide range of broadleaf weeds in fallow, lucerne, maize, millets, pastures, sorghum, sugar cane, sweet corn, winter cereals. Also for the control of woody weeds in agricultural non-crop areas, commercial and industrial areas, pastures and rights-of-way as specified in the Directions For Use Table

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE
OPENING OR USING THIS PRODUCT**

**CONTENTS: 20L
(1L, 5L, 110L, 200L, 1000L)**

APVMA Approval No 70149/62623

Wellfarm Pty Ltd
ACN 158 518 551
22 Calypso Crescent
Point Cook Vic 3030
www.wellfarm.com.au



DIRECTIONS FOR USE

RESTRAINTS

- DO NOT apply to plants that may be stressed (NOT actively growing) due to prolonged periods of extreme cold, moisture stress (water-logged or drought affected) poor nutrition, presence of disease, or previous herbicide treatment as reduced levels of control may result.
- Thorough coverage of both foliage and stems, to the point of runoff, is essential for high volume applications (see GENERAL INSTRUCTIONS; application methods WOODY WEED SITUATIONS section).
- DO NOT spray if rain is likely to occur within 1 hour.

Table 1: Woody Weeds in Agricultural Non-Crop Areas and Rights-of-Way, Commercial and Industrial Areas, Forests and Pastures.

- Legumes present at the time of spraying will be severely damaged.

HIGH VOLUME APPLICATION: Dilute product with water. See General Instructions – Application Method for application details				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE mL/ 100L WATER	CRITICAL COMMENTS
Bathurst burr, Noogoora burr	Seedlings and young plants up to 40cm high	NSW, NT, QLD, WA only	75	
Black bindweed (Climbing buckwheat)	Seedlings and young plants before flowering	NSW, QLD only	300	
Mimosa pigra	Apply from mid to late summer	NT, WA only		
Common sensitive plant	Seedlings and young plants up to flowering	QLD, WA only	500	Add Wellfarm Infuse Spray Adjuvant (see General Instructions; Oils and surfactants).
Bellyache bush		QLD, NSW, WA only		
Blackberry nightshade, Bokhara clover		NSW, QLD only		
Caltrop (yellow vine) (Tribulus terrestris) (T. micrococcus)				
Cobblers pegs	Up to 15cm high			
Cockspur thorn	Up to 3m high			
Creeping lantana	At flowering			
Crofton weed, Mistflower	Seedlings and young plants up to flowering			
Docks (Rumex spp.)	Seedlings and rosettes up to 30cm high			

Hexham scent	Seedlings and young plants up to flowering			Boom Spray: Wellfarm Fluroxypyr 200 EC Herbicide at 0.3L/ha + 0.5L/ha of 2,4-D amine (500g/L).
Honey locust	Seedlings and young plants up to 2m high			
Small flowered mallow (Marshmallow) (Malva parviflora)	Seedlings and young plants up to flowering			
Yellowflower Devil's claw	Seedlings and young plants up to flowering			

Table 1: Woody Weeds in Agricultural Non-Crop Areas and Rights-of-Way, Commercial and Industrial Areas, Forests and Pastures.

HIGH VOLUME APPLICATION: Dilute product with water. See General Instructions – Application Method for application details				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE mL/ 100 L WATER	CRITICAL COMMENTS
Lantana	Seedlings and regrowth 0.5 to 1.2 m high	NSW, QLD only	500	Apply to actively growing plants from October to April. Some regrowth may occur particularly when treating old woody plants with sparse canopies.
	Plants and regrowth 1.2 to 2 m high		1000	
Blue heliotrope	Flowering			
Limebush	Infestations up to 1.5m high only			
Madeira vine	Apply at time of active growth		500	
Milkweed (Euphorbia heterophylla)	3 leaf to flowering	QLD only	1000	Repeat applications will be necessary to control subsequent germinations.
Common sowthistle	Seedlings and young plants up to bolting	NSW, QLD only	500	Add a surfactant (see GENERAL INSTRUCTIONS; Oils and surfactants).
Mother-of-millions (Kalanchoe spp.)	Seedling and young plants before flowering		600	
Prickly acacia	Seedling and young plants up to 2m high	QLD only	750	Add Infuse* Spraying Oil (see GENERAL INSTRUCTIONS; Oils and surfactants). Consult Tropical Weeds Research Centre, Charters Towers, for specific advice on application.

Sida spp.	Seedling and young plants up to flowering	NSW, NT, QLD, WA	1000	
Broadleaf Pepper tree (Schinus terebinthifolius)	Mature leaves, fruiting	QLD only	500	Winter application only. Contact Alan Fletcher Research Station for more information.
Flannel weed (Sida cordifolia)				
Snakeweed (Dark and light blue)	Seedling and young plants before flowering		750	Add Infuse* Spraying Oil (see GENERAL INSTRUCTIONS; Oils and surfactants).
Stinking Passion Flower	Established plants and regrowth	QLD, NT, WA	450	Use 70mL/15 L for a knapsack.
Wandering jew (Tradescantia albiflora)	Young plants up to and including flowering	All States	1500	Some regrowth will usually occur and will require retreatment.
Wattles (including Acacia aulacocarpa A. decora A. harpophylla A. leiocalyx A. salicina)	Seeding plants or regrowth 0.5 to 1.2 m high	NSW, QLD only	500	Apply to actively growing plants when soil moisture is plentiful. Some regrowth may occur particularly when treating old woody plants with sparse canopies and under dry conditions.
	Plants or regrowth 1.2 to 2.0 m high only		1000	

Table 1: Woody Weeds in Agricultural Non-Crop Areas and Rights-of-Way, Commercial and Industrial Areas, Forests and Pastures.

BASAL BARK AND CUT STUMP APPLICATION: Dilute product with diesel. See General Instructions – Application Method for application details				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE L/100 L of diesel	CRITICAL COMMENTS
Celtis (Celtis sinensis)	Basal Bark only: Young plants up to 2m high and 20cm basal diameter	QLD only	3.5	Treat stems from ground level to where multi-stemmed trunks branch.
Chinee apple	Up to 15cm basal diameter		3	With basal bark, treat circumference of stem to a height of 45cm from the ground. Contact the Land Protection Branch, Department of Lands, QLD, for further information on Chinee Apple.
Cockspur thorn	Basal Bark only: Up to 5cm basal diameter		2	
Mimosa bush (Acacia farnesiana)	Up to 5cm basal diameter	QLD, WA only	3	

Prickly acacia	Up to 10cm basal diameter	QLD only	1.5	
Honey locust	Plants up to 10cm basal diameter	QLD, NSW only	1.5	With basal bark, treat circumference of stem to a height of 45cm from the ground. For cut stump application use a rate of 5L/100 diesel for all plant sizes. Contact the Land Protection Branch, Department of Lands, QLD, for further information on Honey Locust.
	Plants 10 to 20 cm basal diameter		3	
	Plants >20cm basal diameter		5	
Sisal hemp (Agave spp.)	All growth stages	QLD only	3	Treat as an overall spray. Contact The Land Protection Branch, Department of Lands, QLD for advice to control large infestations.
			10mL undiluted product per plant	Lever out centre of plant with crowbar and immediately treat the exposed cut area.

BROADCAST AND AERIAL APPLICATION: Dilute product with water. See General Instructions – Application Method for application details				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE L/ha	CRITICAL COMMENTS
Mimosa pigra	Actively growing plants	NT, WA only	3L	Aerial application: Add Wellfarm Infuse Spray Adjuvant at the rate of 1 L/100 L spray mix. Apply to actively growing plants from mid to late summer. Contact the Department of Primary Industries and Fisheries, NT for further information.

LOW VOLUME, HIGH CONCENTRATE APPLICATION: Use a drench gun or gas-powered gun. See General Instructions – Application Method for application details				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE Per 10 L of water	CRITICAL COMMENTS
Limebush	Isolated bushes up to 1.2m high only	NSW, QLD only	1L	Apply a 50mL dose per 5m ² of bush surface area.
Tree violet (Hymenanchera dentata)	Apply from late flowering to green fruit up to 1.2m high	NSW only		Apply a 50mL dose per cubic metre of bush.

Table 2: Established Grass Pastures

WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE L/ha	CRITICAL COMMENTS
Blue billygoat weed, Common sensitive plant, Giant sensitive plant, Spinyhead sida	Apply before flowering	QLD, WA only	1.5	Add Wellfarm Infuse Spray Adjuvant at 1 L/ha
St John's wort	Apply from bud to full bloom (usually late Nov to early Jan)	ACT, NSW and VIC only	3	Some regrowth will occur. Treat regrowth the following season for best results. Use at least 200L water/ha.
Silverleaf nightshade	From onset of flowering to early berry-set (usually spring to mid-summer)	NSW only	0.75 or 0.375 + 1.5 - 2 2,4-D amine (500g/L)	Add Wellfarm Infuse Spray Adjuvant at 1L/ha. To ensure maximum effect, delay application until the majority of shoots have emerged. Follow-up treatment of regrowth is critical for best control.

Table 3: Sorghum, Maize, Millets and Sweet Corn (NSW & QLD Only)

CROP	CROP GROWTH STAGE	WEEDS CONTROLLED	WEED GROWTH STAGE	RATE L/ha	CRITICAL COMMENTS
Sorghum	Apply when secondary roots are present, from 4 fully expanded leaves (15cm tall) up to boot (also see CRITICAL COMMENTS)	Annual ground cherry, Wild gooseberry (Physalis spp.)	2 to 8 leaf Up to 15cm tall	0.5	Sorghum: From 8 leaf to boot stage, use dropper nozzles to prevent herbicide coming in contact with the crop's leaves and the growing point (meristem).
			15 to 30 cm tall	0.75	
		Apple-of-Peru	Seedling plants up to 15cm tall		
		Bathurst burr, Noogoora burr	2 to 8 leaf Up to 20cm tall	0.5	
Maize & Sweet corn	Apply when secondary roots are present, from 3 fully expanded leaves (10cm tall) up to just before tasselling (see CRITICAL COMMENTS)		20 to 50 cm tall	0.75	Maize and Sweet Corn: From 6 leaf to just before tasselling, use dropper nozzles to prevent the herbicides coming in contact with the crop's leaves and the growing point (meristem).
		Pigweed (Portulaca oleracea)	Up to 10cm diameter	0.5	
			10 to 30 cm diameter	0.75	
		Sesbania pea	2 to 6 leaf Up to 10cm tall	1.5	
Millets	Spray when secondary roots have developed, usually early to mid-tillering, and NOT later than before heads start to form at the base of tillers. (See CRITICAL COMMENTS)	Silverleaf nightshade (NSW only) ⁽¹⁾	Full flower to early berry	0.75 + Infuse at 1L/ha	Millets: DO NOT use mixes with atrazine. ⁽¹⁾ This treatment may be slightly damaging to the crop. To minimise crop damage apply using dropper nozzles at all crop stages.
		Starburr (Acanthospermum hispidum) (QLD only)	Up to 12 leaf and before flowering	1.5 or 0.75 + 2 L atrazine (500g/L)	
		Thornapples (Datura spp.)	2 to 8 leaf Up to 15cm tall	0.75	
		Volunteer sunflower	2 to 5 leaf Up to 20cm tall	1	

Table 3: Sorghum, Maize, Millets and Sweet Corn (NSW & QLD Only)

Wellfarm Fluroxypyr 200 EC Herbicide in tank-mixes with atrazine: Sorghum, Maize and Sweet Corn.					
CROP	CROP GROWTH STAGE	WEEDS CONTROLLED	WEED GROWTH STAGE	RATE L/ha	CRITICAL COMMENTS
Sorghum, Maize & Sweetcorn	Spray when secondary roots have developed, usually early to mid-tillering and NOT later than before heads start to form at the base of the tillers (See CRITICAL COMMENTS)	Amaranthus spp. Including: Boggabri weed, Dwarf amaranth, Green amaranth, Redshank, Anoda weed, Bladder ketmia, Black pigweed (Trianthema portulacastrum), Caltrop (yellow vine), including Tribulus terrestris, T. microccus and T. maximus Cowvine (peach vine) (Ipomoea lonchophylla), Hairy wandering jew (Commelina benghalensis), Mintweed	Seedling plants up to 15cm tall or rosettes up to 15cm diameter	0.5 + 1.5 of atrazine flowable (500g/L) or 0.75 + 2 of atrazine flowable (500g/L)	Use the low rate (0.5 + 1.5 L) when weeds are small (5-7 cm tall/ diameter). Use the high rate (0.75 + 2 L) when the weeds are larger (7 - 15 cm tall/ diameter). Wellfarm Fluroxypyr 200 EC Herbicide is generally more compatible with Liquid Atrazine Products (see GENERAL INSTRUCTIONS; compatibility section). Add a surfactant (See GENERAL INSTRUCTIONS; Oils and Surfactants). DO NOT add an oil to mixtures of Wellfarm Fluroxypyr 200 EC Herbicide and Atrazine.
		Euphorbia davidii	Cotyledons to 4 nodes up to 15 cm	1 + 2 atrazine flowable (500 g/L)	
		Volunteer peanuts	Up to 15 cm diameter	1 + 4.5 atrazine flowable (500 g/L)	

Sweet Corn: TAS Only					
CROP	CROP GROWTH STAGE	WEEDS CONTROLLED	WEED GROWTH STAGE	RATE L/ha	CRITICAL COMMENTS
Sweet corn only	3 to 5 leaf	Blackberry nightshade, Volunteer potatoes	3 to 5 leaf	1	

Table 4: Winter Cereals (Wheat, Barley, Oats and Triticale)

CROP GROWTH STAGE	WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE L/ha	CRITICAL COMMENTS
Apply from 3 leaf to flag (Zadoks 13 to 39)	Bedstraw (Galium tricornutum)	1 to 3 whorl	VIC, SA, WA	1	⁽¹⁾ Add either Infuse or a surfactant (see GENERAL INSTRUCTIONS: Oils and Surfactants).
	Cleavers (Galium aparine)		NSW, VIC only		
	Black bindweed (Climbing buckwheat)	2 to 4 leaf	NSW, QLD only	0.5 ⁽¹⁾	Useful suppression only.
		2 to 6 leaf		0.75 or 0.5 + 5 g Metsulfuron methyl ⁽¹⁾	
	Common sowthistle (Sonchus oleraceus)	2 to 5 leaf		1	
	Deadnettle	2 to 6 leaf		1.5 or 0.5 + 5 g Metsulfuron methyl ⁽¹⁾	
	Spiny emex (Doublegee, Three cornered jack)	2 to 4 leaf	NSW, SA, QLD, WA		
	Prickly lettuce	2 to 5 leaf	NSW, QLD, TAS, VIC, WA	1	
	Volunteer lupins	2 to 8 leaf	NSW, VIC, WA only	1.5	
	Volunteer potato	10 to 15 cm tall	WA and TAS only		Plants 15 to 30 cm tall will only be suppressed.
	Wireweed	2 to 3 leaf	NSW, QLD, SA, TAS, VIC and WA		
			NSW and QLD only	0.5 + 5 g Metsulfuron methyl ⁽¹⁾	

	Bittercress (<i>Coronopus didymus</i>), Mustards, Shepherd's purse, Turnip weed, Wild radish, Wild turnip	Up to 8 leaf and up to 15 cm diameter	QLD, NSW, VIC, SA, TAS, WA	0.5 to 1.5 + Metsulfuron methyl ⁽¹⁾ or Eclipse ⁽¹⁾ or MCPA LVE or MCPA amine	The Wellfarm Fluroxypyr 200 EC Herbicide rate depends on what other weeds are present as listed above. See Mixtures comment above. Metsulfuron methyl (600g/kg) @ 5 g/ha (this mix does not control wild radish). Eclipse @ 5-7 g/ha (use the 5 g rate on turnip weed only). MCPA LVE (500g/L) @ 700 mL/ha. MCPA Amine (500g/L) @ 1.0 L/ha.
--	---	---	--	--	--

Table 5: Summer Fallow

WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE L/ha	CRITICAL COMMENTS
Annual ground cherry, Wild gooseberry (<i>Physalis</i> spp.)	2 to 8 leaf, up to 15cm tall	NSW, QLD only	0.75 ⁽²⁾	⁽¹⁾ Add Infuse* Spraying Oil (see GENERAL INSTRUCTIONS; Oils and Surfactants). When mixing with Glyphosate 450g/L to control both grass and broadleaf weeds, refer to the Glyphosate 450g/L label for use rates and adjuvants recommended for the grasses (see GENERAL INSTRUCTIONS; compatibility section).
Bathurst burr, Noogoora burr	2 to 8 leaf, up to 20cm tall	NSW, QLD, VIC, WA only		
Bellvine	Pre-flowering	NSW, QLD only	0.5 + 1.2 Glyphosate 450g/L	⁽²⁾ Delay treatment until the maximum number of shoots have emerged, but before the onset of fruiting (late summer). DO NOT treat plants showing symptoms from previous treatment. Use the high rate when longer-term weed control (6-10 months) is required and delay planting crops during this period. The low rate will require follow- up treatments.
Bladder ketmia	4 to 8 leaf, up to 10cm tall			
Cowvine (Peach vine) <i>Ipomoea lonchophylla</i>	2 to 10 leaf up to 10cm diameter			
Caltrop (yellow vine), including <i>Tribulus terrestris</i> , <i>T. maximus</i> and <i>T.</i> <i>microccus</i>	Up to 15cm diameter		0.5 + 1.0 Glyphosate 450g/L	
Pigweed (<i>Portulaca oleracea</i>)	Up to 10cm diameter		0.75 ⁽¹⁾	
	Up to 60cm diameter		0.75 + 1.0 Glyphosate 450g/L	
<i>Polymeria pusilla</i>	2 to 10 leaf up to 20cm diameter		1 ⁽¹⁾ or 0.5 + 1.2 Glyphosate 450g/L	

WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE L/ha	CRITICAL COMMENTS
Rhynchosia	Seedlings to early flowering		1 ⁽¹⁾ or 0.375 + 0.8 Glyphosate 450g/L	
Smallflower mallow or Marshmallow (Malva parviflora)	Up to 8 leaf up to 20cm diameter		1 ⁽¹⁾	
Thornapples (Datura spp.)	2 to 8 leaf up to 15cm diameter	NSW, QLD, WA only	0.75 ⁽¹⁾ or 0.5 + 1.2 Glyphosate 450g/L	
Sesbania pea	2 to 6 leaf up to 10cm tall	NSW, QLD only	1.5 ⁽¹⁾ or 0.5 + 1.2 Glyphosate 450g/L	
Perennial Ground Cherry (Physalis virginiana) ^(w)	Bud to early flowering up to 20cm tall		1.5 or 3 ⁽¹⁾	
Silverleaf nightshade	Full flower to early berry-set (usually Dec – Feb)	NSW only	0.75 or 0.375 + 1.5 – 2 2,4-D amine (500g/L)	Add Wellfarm Infuse Spray Adjuvant at the rate of 1 L/100L spray mixture. To ensure maximum effect, delay application until the majority of shoots have emerged. Follow-up treatment will be required to control regrowth and is critical for optimum control. If wanting to prevent seed set repeat applications may be needed in the same season, although this does NOT lead to better long-term control.
Volunteer peanuts	Up to 15cm diameter	QLD only	1 + 4.5 atrazine flowable (500g/L)	Add a surfactant (see General Instructions; Oils and Surfactants). Important: (see GENERAL INSTRUCTIONS; Compatibility Section).
Volunteer sunflowers	2 to 5 leaf up to 20cm	NSW, QLD only	1	Add Wellfarm Infuse Spray Adjuvant (see General Instructions; Oils and surfactants section).

Table 6: Winter Fallow

WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE L/ha	CRITICAL COMMENTS
Bedstraw (Galium tricornutum)	Up to 5 whorl	VIC, SA, WA only	1 ⁽¹⁾	(1) Add Wellfarm Infuse Spray Adjuvant (see GENERAL INSTRUCTIONS; Oils and Surfactants Section). (2) Add Infuse or a surfactant (see GENERAL INSTRUCTIONS; Oils and Surfactants Section).
Cleavers (Galium aparine)		NSW, VIC only		
Black bindweed (Climbing buckwheat)	2 to 8 leaf up to 10cm diameter	NSW, QLD only	0.75 ⁽¹⁾	When mixing with Glyphosate 450g/L to control both grass and broadleaf weeds, refer to the Glyphosate 450g/L label for use rates and adjuvants recommended for the grasses (see GENERAL INSTRUCTIONS; Compatibility Section).
Common sowthistle (Sonchus oleraceus)	2 to 5 leaf up to 10cm diameter		1.0 ⁽¹⁾ or 0.5 + 0.6 Glyphosate 450g/L	
Prickly lettuce				
Spiny emex (Doublegee, Three cornered jack)	2 to 8 leaf		1.5 ⁽¹⁾ or 0.5 ⁽²⁾ + 5 g Metsulfuron methyl (600g/kg)	
Wireweed	2 to 3 leaf up to 10cm tall		1.5 ⁽¹⁾ or 0.5 ⁽²⁾ + 5 g Metsulfuron methyl (600g/kg) or 0.5 ⁽²⁾ + 0.6 Glyphosate 450g/L	

Table 7: Sugar Cane (QLD, NSW, NT and WA Only)

CROP STAGE GROWTH	WEEDS CONTROLLED	WEED GROWTH STAGE	RATE L/ha	CRITICAL COMMENTS
From early tillering to maturity	Balsum pear, Blackberry nightshade, Blue billygoat weed, Centro, Cowpea, Giant sensitive plant, Lablab bean, Noogoora burr, Phasey bean, Pinkburr, Prickly African Cucumber, Spinyhead sida, Stinking passion flower (seedlings only)	Apply from 2 to 3 leaf until flowering	Ground: 1.3 Aerial: 1.5	For optimal weed control, delay application until just before the “close-in” stage. Aerial Application: Apply in NOT less than 60L/ha water and add Wellfarm Infuse Spray Adjuvant at 1L/100L spray mixture. Ground Application: Apply in 100 – 400 L/ha water and add Wellfarm Infuse Spray Adjuvant at 500mL/100L of spray mixture.
	Bellvine, Morning glory, Red or pink convolvulus, Star-of-Bethlehem		As above + 1 2,4-D amine (500g/L)	
	Stinking passion flower	Established or ratoon plants with at least 1.0m of regrowth	High Volume: 450mL/100L water Knapsack 70mL/15L water	Thoroughly wet plants to the point of run-off.
	Milkweed (Euphorbia heterophylla)	Seedlings and young plants up to flowering.	3 or 2.3 + 4 atrazine flowable (500g/L)	Better control will be achieved with the atrazine mixture. Delay application until just before the cane reaches the “close-in” stage. This will improve control and minimise the number of seedlings that germinate.

Table 8: Lucerne (NSW Only)

CROP STAGE GROWTH	WEEDS CONTROLLED	WEED GROWTH STAGE	RATE L/ha	CRITICAL COMMENTS
Established crops at least 18 months old	Annual ground cherry, Bathurst burr, Noogoora burr, Wild gooseberry	2 to 8 leaf up to 15cm high	0.5	<p>To minimise crop injury and to maximise weed control, cut, slash or heavily graze the lucerne before application. Wherever possible, irrigate before application to stimulate weed growth.</p> <p>DO NOT treat crops growing on sandy or stony soils.</p> <p>DO NOT treat crops after the summer growing season (after end of March).</p> <p>To broaden the spectrum of weeds controlled, Wellfarm Fluroxypyr 200 EC Herbicide can be mixed with 2,4-DB Amine.</p>
	Pigweed	Up to 10cm diameter		

Table 9: Poppies (TAS Only)

CROP STAGE GROWTH	WEEDS CONTROLLED	WEED GROWTH STAGE	RATE L/ha	CRITICAL COMMENTS
4 to 6 leaf	Cleavers, Fumitory	2 to 6 leaf	1	
	Shepherd's purse, Wireweed		1 + 5 Asulox	
8 to 10 leaf	Common sowthistle, Prickly lettuce	2 to 5 leaf	1	DO NOT apply Wellfarm Fluroxypyr 200 EC Herbicide to poppies later than the 8 to 10 leaf growth stage as a reduction of alkaloid content could occur.
	Black nightshade	Cotyledon to 4 leaf	1.5	
	Fumitory	6 to 10 leaf		
	Volunteer potato	From tuber initiation to flower bud		

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

WITHHOLDING PERIODS:

CROPS AND PASTURES	DO NOT GRAZE FAILED CROPS AND TREATED PASTURES OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.
POPPIES	DO NOT SPRAY POPPIES LATER THAN 10 WEEKS BEFORE HARVEST.

MINIMUM RE-CROPPING PERIODS

PLANT-BACK PERIODS FOR CROPS FOLLOWING THE APPLICATION OF WELLFARM FLUROXYPYR 200 EC HERBICIDE FOR RATES UP TO 1.5 L/HA			
RATE L/ha	0.375	0.75	1.5
CROP	DAYS		
Barley	7	7	7
Wheat	7	7	7
Chickpea	7	7	7
Cotton	14	14	28
Soybean	7	7	14
Sunflower	7	7	7
Maize	7	7	7
Sorghum	7	7	7

NOTE: Before using Wellfarm Fluroxypyr 200 EC Herbicide in tank mixes with other herbicides, check the plant-back information on all product labels. The time between spraying and planting will be determined by the most residual product, i.e. the product with the longest plant-back period.

GENERAL INSTRUCTIONS

MIXING

Wellfarm Fluroxypyr 200 EC Herbicide may be mixed with water or diesel.

Mix only sufficient chemical for each day's use and avoid storing.

Mixing in Water: Half fill the spray tank with water and add the required quantity of Wellfarm Fluroxypyr 200 EC Herbicide and complete filling. Agitate continuously to ensure thorough mixing before and during application.

Mixing in Diesel: Half fill the tank with diesel and add the required quantity of Wellfarm Fluroxypyr 200 EC Herbicide. Add the remainder of the diesel and agitate or shake to mix contents.

Tank Mixtures: Wettable powder or dry flowable formulations (e.g. water dispersible granules) should be added to the spray tank first, followed by suspension concentrates (flowables), water soluble salts and then emulsifiable concentrate formulations (Wellfarm Fluroxypyr 200 EC Herbicide). Add spraying oils and surfactants (wetters) last.

OILS AND SURFACTANTS

Oils

Use only Wellfarm Infuse Spray Adjuvant at the rate of 500mL/100L of spray mix. When using less than 100L/ha spray volume, ensure a minimum of 250mL/ha of Infuse is used, unless 1L/100L or 1L/ha is specified.

Surfactants (wetters)

Use a 1000g/L non-ionic surfactant such as Wellfarm Cropwett 1000 at 100mL/100L of spray mix where required.

COMPATIBILITY

Wellfarm Fluroxypyr 200 EC Herbicide Is Compatible With The Herbicides Listed. Follow any regional restrictions, and all directions and restrictions on the label, of any chemical mixed with Wellfarm Fluroxypyr 200 EC Herbicide.

Atrazine (see below)

Metsulfuron-Methyl (600g/kg)

Broadstrike

Eclipse

Diclofop-Methyl

Triclopyr (600 g/L)

Clopyralid

MCPA

Glyphosate 360g/L

Glyphosate 450g/L

Clodinafop 240 EC (see below)

Picloram + 2,4-D

Picloram + MCPA

2,4-D

2,4-DB

ATRAZINE

AVOID USING HARD WATER WHEREVER POSSIBLE.

Where hard water CANNOT be avoided, the addition of CALGON water conditioning agent to the spray tank, at 100g/100L water, before adding any herbicide may improve compatibility.

AGITATION IS VERY IMPORTANT WHEN MIXING WELLFARM FLUROXYPYR 200 EC HERBICIDE AND ATRAZINE.

Wellfarm Fluroxypyr 200 EC Herbicide plus Atrazine tank mixes must be agitated vigorously and continuously during mixing and application. After mixing DO NOT allow to stand without agitation. Ensure that the time from mixing to the end of application is NOT more than 2 hours. If settling out occurs re-suspension is difficult, even with vigorous agitation.

Agitation using only the pump's by-pass is usually inadequate, particularly with larger tanks (more than 2000L). Additional mechanical agitation will be necessary in large tanks, computer sprayers and mixing tanks.

When additional surfactant is required, add a 100% concentrate non-ionic surfactant at 100mL/100L of spray mix. DO NOT use a spraying oil when tank mixing Wellfarm Fluroxypyr 200 EC Herbicide and Atrazine.

Guidelines For Tank-Mixing Wellfarm Fluroxypyr 200 EC Herbicide and Common Atrazine Formulations

Tank Mix	Rate (L/ha)	Water Hardness			Minimum Water Volume (L/ha)		Critical Comments
		Soft	Medium	Hard	Ground	Aerial	
Wellfarm Fluroxypyr 200 EC Herbicide	0.75	✓	✓	✓	50	35	
Wellfarm Fluroxypyr 200 EC Herbicide + Gesaprim 500 FW	0.75 + 2	✓	✓	✓	50 – 100	35	Precipitate can be easily resuspended.
Wellfarm Fluroxypyr 200 EC Herbicide + Atradox 900 WG	0.75 + 1.1	✓	✗	✗	100	DO NOT use	Precipitate may be difficult to resuspend and may block nozzles.
Wellfarm Fluroxypyr 200 EC Herbicide + Nu-Trazine DF	0.75 + 1.1	✓	✗	✗	100	DO NOT use	Sediment may be difficult to resuspend and may block nozzles.
Wellfarm Fluroxypyr 200 EC Herbicide + Nu-Trazine 500 FW	0.75 + 2	✓	✓	✗	100	DO NOT use	Precipitate may be difficult to resuspend and may block nozzles.

Wellfarm Clodinafop 240 EC Herbicide

Always use Wellfarm Infuse Spray Adjuvant with Wellfarm Fluroxypyr 200 EC Herbicide + Wellfarm Clodinafop 240 EC Herbicide tank-mixes at 500mL/100L of spray mix with a minimum of 250mL/ha. DO NOT mix Wellfarm Fluroxypyr 200 EC Herbicide with Wellfarm Clodinafop 240 EC Herbicide if the grass weeds are NOT actively growing. Always use the maximum label rate of Wellfarm Clodinafop 240 EC Herbicide for the appropriate grass growth stage.

DO NOT use Wellfarm Fluroxypyr 200 EC Herbicide at more than 0.75L/ha in tank mixes with Wellfarm Clodinafop 240 EC Herbicide.

Glyphosate 450g/L

When mixing Wellfarm Fluroxypyr 200 EC Herbicide with Glyphosate 450g/L to control both grass and broadleaf weeds, refer to the Glyphosate 450g/L label for use rates and adjuvants recommended for the grasses. DO NOT use Glyphosate 450g/L at less than 1.2L/ha in tank mixes with Wellfarm Fluroxypyr 200 EC Herbicide, when barnyard grass, buttongrass, crowsfoot grass, native millet and liverseed grass are the target species.

APPLICATION METHODS AND WATER RATES

BROADCAST APPLICATION IN CROPPING, PASTURE AND FALLOW SITUATIONS

A. Ground Application (Boom)

Apply Wellfarm Fluroxypyr 200 EC Herbicide with an accurately calibrated boom sprayer, in at least 50L/ha water (100-400 L/ha for sugar cane).

Flat fan nozzles are recommended using pressures in the range 200 to 300 kPa.

Set the boom at a height to ensure a double overlap of the nozzle patterns.

B. Ground Directed Application (Dropper Nozzles)

To minimise crop effects, dropper nozzles should be used in sorghum when the crop is beyond the 8 leaf growth stage and in maize and sweet corn when the crop is beyond the 6 leaf growth stage.

Adjust the nozzles to direct the spray into the base of the crop and away from the leaves and the growing point. See manufacturer's directions for setting up and calibration of dropper nozzles.

C. Aerial Application

Apply in a minimum volume of at least 35L/ha water (60L/ha in sugarcane).

Use equipment calibrated to produce droplets with an average diameter (Volume Mean Diameter; VMD) of 250 – 350 micron.

DO NOT apply when the temperature is above 30°C, when there is NO wind or when the wind is blowing toward susceptible crops.

WOODY WEED SITUATIONS

Weeds must be actively growing to attain optimal effect. Delay the treatment of regrowth following bulldozing, slashing, burning, ploughing or a previous chemical treatment until it has at least 1 metre of new, vigorous, growth.

A. High Volume Application

Hand Gun

Apply the recommended mix to obtain full coverage of leaves and stems using a number 6 – 8 tip at 700 to 1500 kPa. To obtain good coverage, a spray volume of 1500 to 4000 L/ha (15 to 40 L/100m²) is required per infested hectare.

Ensure thorough coverage to the point of runoff.

Knapsack

Knapsack sprayers may be used on smaller infestations where penetration and coverage of the canopy is easier to achieve. Use the same use rate and spray techniques as for handgun application.

B. Low Volume, High Concentrate Application

Drench Gun or Gas-Powered Gun

Apply the recommended mixture uniformly across the foliage by applying 50mL shots to cover 4 to 5 m² of surface area of plant. This is approximately equivalent to 20 droplets per cm² of the leaf surface.

Use a marking agent as recommended by the equipment manufacturer to check spray coverage.

C. Basal Bark and Cut Stump Application

Basal Bark

DO NOT apply to wet stems as this can repel the diesel mixture.

Spray or paint the recommended mixture around the base of each stem from ground level to a height of at least 30cm from the ground, wetting the bark to the point of runoff.

Apply with a paint brush or a pressure sprayer with an approximate lance and solid cone nozzle. If using spray equipment use low pressures (</_ 200 kPa) sufficient to form a cone of spray.

Old rough bark will require more spray than smooth or young thin bark.

Cut Stump

Apply the recommended mixture liberally to the freshly cut stump immediately after cutting.

Apply by spraying or painting the cut surface and sides of the stump.

Best results are obtained when the stems are cut less than 15cm above the ground.

CLEANING SPRAY EQUIPMENT

Rinse Water Should Be Discharged Onto A Designated Disposal Area Or, If This Is Unavailable, Onto Wasteland Away From Desirable Plants And Water-Courses.

Cleaning Equipment After Using Water-Based Sprays:

Rinsing: After using Wellfarm Fluroxypyr 200 EC Herbicide, empty the tank completely and drain the whole system. Thoroughly wash inside the spray unit using a pressure hose. Drain and clean any filters in the tank, pump, lines, hoses and nozzles.

After cleaning the tank as above, quarter fill the clean water and circulate through the pump, lines and nozzles. Drain and repeat the rinsing procedure twice.

Decontamination (Before Spraying Cotton And Other Sensitive Crops; See Protection Of Crops): Wash the tank and rinse the system as above. Then quarter fill the tank and add an alkali detergent (e.g. liquid SURF, OMO, DRIVE) at 500mL/100L of water or the powder equivalent at 500g/100L and circulate throughout the system for at least 15 minutes.

Drain the whole system. Remove filters and nozzles and clean them separately. Finally flush the system with clean water and allow to drain.

Cleaning Equipment After Using Diesel - Based Sprays:

On completion of spraying, use a degreaser such as Caltex Kwik-D-Grease to remove traces of diesel from the sprayer. Rinse tank and spray through nozzles with water to remove degreaser.

Then quarter fill the tank and add an alkali detergent (e.g. liquid SURF, OMO, DRIVE) at 50mL/10L of water or the powder equivalent at 50g/10L of water. Shake sprayer to circulate the washing solution throughout the sprayer, then spray the solution through the nozzles. Rinse well with clean water to remove the detergent.

To clean brushes and containers, spray liberally with degreaser. Hose off with clean water and repeat using detergents as above.

DO NOT use this equipment for any other purpose.

RESISTANT WEEDS WARNING

GROUP	I	HERBICIDE
--------------	----------	------------------

Wellfarm Fluroxypyr 200 EC Herbicide is a member of the pyridine group of herbicides. Wellfarm Fluroxypyr 200 EC Herbicide has the disrupters of plant cell growth mode of action. For weed resistance management, Wellfarm Fluroxypyr 200 EC Herbicide is a Group I herbicide.

Some naturally occurring weed biotypes resistant to Wellfarm Fluroxypyr 200 EC Herbicide and other Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if Wellfarm Fluroxypyr 200 EC Herbicide and other Group I herbicides are used repeatedly. These resistant weeds will NOT be controlled by Wellfarm Fluroxypyr 200 EC Herbicide or any other Group I herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Wellfarm Pty Ltd accepts NO liability for any losses that may result from the failure of Wellfarm Fluroxypyr 200 EC Herbicide to control resistant weeds.

Wellfarm Fluroxypyr 200 EC Herbicide may be subject to specific resistance management strategies. For further information consult your local farm chemical supplier, consultant, local Department of Agriculture or Primary Industries.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Susceptible crops include but are NOT limited to clovers, cotton, fruit, hops, lupins, ornamentals, peas, pine tree, potatoes, navy beans, safflower, shade trees, soybeans, sunflower, tobacco, tomatoes, vegetables and vines.

Wellfarm Fluroxypyr 200 EC Herbicide can be damaging to susceptible crops during both growing and dormant periods.

Grasses are normally unaffected by Wellfarm Fluroxypyr 200 EC Herbicide and establish quickly after treatment. Transitory damage can occur on some species particularly those that spread by stolons such as couch grass (*Cynodon dactylon*), Kikuyu grass and carpet grass (*Axonopus* sp.)

DO NOT allow spray to drift onto susceptible crops, shade trees and *Pinus* spp.

DO NOT use under weather conditions or from spraying equipment that could cause spray to drift onto nearby susceptible plants.

PROTECTION OF LIVESTOCK

DO NOT graze or cut treated crops or plants for food except as specified under withholding periods.

Poisonous plants may become more palatable after spraying therefore stock should be kept out of the area until the plants have died down.

DO NOT allow stock to re-enter paddocks containing treated poisonous plants, until the plants have died down.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate dams, streams, drains, rivers or waterways with the chemicals or used containers.

Alongside waterways, treat only noxious weeds and poisonous plants.

STORAGE AND DISPOSAL

KEEP OUT OF REACH OF CHILDREN

Storage For All Containers

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight.

Disposal Recycled Containers

This container can be recycled if it is clean, dry, free of visible residues and has the drum MUSTER logo visible. Triple or preferably pressure rinse containers before disposal. Dispose of rinse by adding to the spray tank. DO NOT dispose of undiluted chemicals on site. Wash outside of the container and cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at a drum MUSTER collection or similar container management site. The cap should NOT be replaced but taken separately.

Disposal Non-Recycled Containers

Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty containers for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty containers 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory Government Regulations. DO NOT burn empty containers or product.

Disposal Refillable Containers

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SMALL SPILL MANAGEMENT

Wear protective equipment (see SAFETY DIRECTIONS). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal (see STORAGE AND DISPOSAL section). If necessary wash the spill area with an alkali detergent and water and absorb the wash liquid for disposal.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. Sensitive workers should use protective clothing when opening the container and preparing the spray and using the prepared spray. Wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and face shield or goggles. After each day's use, wash gloves, face shield or goggles and contaminated clothing. Wash hands after use.

FIRST AID

If poisoning occurs contact a Doctor or phone the Australian Poisons Information Centre 13 11 26.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS) that can be obtained from the supplier.

CONDITIONS OF SALE

The use of this product is beyond the control of the manufacturer. No warranty expressed or implied is given by Wellfarm Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Wellfarm Pty Ltd accepts NO responsibility for any consequences whatsoever resulting from the use of this product.

MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY

Wellfarm Pty Ltd has no control over storage, handling and manner of use of this product. Where the material is not stored, handled or used correctly and in accordance with the directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory

warranties) will apply. Wellfarm Pty Ltd accepts NO liability for any losses or damages arising from incorrectly storage, handling or use.

IN AN EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE

APVMA Approval No 70149/62623

Wellfarm Pty Ltd
ACN 158 518 551
22 Calypso Crescent
Point Cook Vic 3030
www.wellfarm.com.au