

POISON KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Haloxyfop 520 EC

Herbicide

**ACTIVE CONSTITUENT: 520g/L HALOXYFOP
PRESENT AS THE HALOXYFOP-R METHYL ESTER**

GROUP A HERBICIDE

For the post-emergent control of a wide range of annual and perennial grass weeds in grain legume and oilseed crops, lucerne, medic and clover pasture and seed crops, forestry, bananas, citrus, grapes, pineapples, pome and stone fruit, pyrethrum, tropical fruit and nut crops as specified in the Directions For Use Table

**IMPORTANT: READ THE ATTACHED LEAFLET
BEFORE OPENING OR USING THIS PRODUCT**

CONTENTS
250ML

STORAGE AND DISPOSAL KEEP OUT OF REACH OF CHILDREN

Store in the closed, original container in a dry, cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT store near feedstuffs, fertilisers or seeds. Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty containers for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty containers 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory Government Regulations. DO NOT burn empty containers or product.

SMALL SPILL MANAGEMENT

Wear protective equipment (see SAFETY DIRECTIONS). Apply absorbent material such as earth, sand, cat litter or clay granules to the spill. When absorption is complete, sweep up material and contain in a refuse vessel for disposal (see STORAGE AND DISPOSAL section). If necessary wash the spill area with an alkali detergent and water and absorb this wash liquid for disposal as directed above.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening container, preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and washable hat, elbow-length PVC gloves and face shield or goggles. After each day's use, wash gloves and face shield or goggles and contaminated clothing. Wash hands after use.

FIRST AID

If poisoning occurs contact a Doctor or phone the Australian

Poisons Information Centre 13 11 26. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a Doctor.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS) that can be obtained from the supplier.

CONDITIONS OF SALE

The use of this product is beyond the control of the manufacturer. No warranty expressed or implied is given by Wellfarm Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Wellfarm Pty Ltd accepts NO responsibility for any consequences whatsoever resulting from the use of this product.

MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY

Wellfarm Pty Ltd has no control over storage, handling and manner of use of this product. Where the material is not stored, handled or used correctly and in accordance with the directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Wellfarm Pty Ltd accepts NO liability for any losses or damages arising from incorrectly storage, handling or use.

APVMA Approval No 70164/0714

**IN AN EMERGENCY
DIAL 000 POLICE
OR FIRE BRIGADE**



Wellfarm Pty Ltd
ACN 158 518 551
ABN 32 158 518 551

22 Calypso Crescent
Point Cook Vic 3030
www.wellfarm.com.au



Batch No

DOM

POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

Wellfarm Haloxyfop 520 EC Herbicide

**ACTIVE CONSTITUENT: 520g/L HALOXYFOP
PRESENT AS THE HALOXYFOP-R METHYL ESTER**

GROUP	A	HERBICIDE
--------------	----------	------------------

For the post-emergent control of a wide range of annual and perennial grass weeds in grain legume and oilseed crops, lucerne, medic and clover pasture and seed crops, forestry, bananas, citrus, grapes, pineapples, pome and stone fruit, pyrethrum, tropical fruit and nut crops as specified in the Directions For Use Table

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE
OPENING OR USING THIS PRODUCT**

**CONTENTS: 250ML
(1L, 5L, 10L, 20L)**

APVMA Approval No 70164/0714

Wellfarm Pty Ltd
ACN 158 518 551
22 Calypso Crescent
Point Cook Vic 3030
www.wellfarm.com.au



DIRECTIONS FOR USE RESTRAINTS

DO NOT apply to weeds which may be stressed (NOT actively growing) due to prolonged periods of extreme cold, moisture stress (waterlogged or drought affected), poor nutrition or previous herbicide treatment as reduced levels of control may result.

DO NOT spray if rain is likely to occur within 1 hour.

Table 1A. Winter Crops – Canola, Chickpeas, Faba Beans, Field Peas, Lentils, Linola, Linseed, Lupins, Lucerne, Vetch, Medic And Clover Pastures Or Seed Crops

Weeds Controlled	Weed Growth Stage	Rate (mL/ha)		Critical Comments
		With Uptake ¹ Spraying Oil	With Non-Ionic Wetter ²	
Annual Ryegrass	2 to 4 leaf	75	100	CANOLA, LINOLA AND LINSEED: DO NOT apply after the 8 leaf stage of the crop. DO NOT apply after the commencement of stem elongation. This means that application must NOT occur after the 8 leaf stage, or if stem elongation commences before the 8 leaf stage, application must NOT occur after stem elongation has commenced. DO NOT apply more than 1 application of herbicide containing haloxyfop per crop. DO NOT apply after grazing.
	early tillering	100	100	
Barley grass Brome grass Paradoxa grass Volunteer cereals	2 to 4 leaf	50	75	^{1 2} See GENERAL INSTRUCTIONS, Spraying oils/wetters section.
	early tillering	75	100	
Wild Oats WA, SA, VIC, TAS, Southern & Central NSW	2 to 4 leaf	37.5	50	FIELD PEAS AND CANOLA: The only oil recommended for use with Wellfarm Haloxyfop 520 EC Herbicide is Uptake* Spraying Oil. Wellfarm Haloxyfop 520 EC Herbicide + Lontrel* 750 SG + Uptake* Spraying Oil are compatible and selective to canola. This tank-mix is also compatible with atrazine and selective to triazine tolerant canola.
	early tillering	50	75	
Wild Oats Northern NSW & QLD	2 to 4 leaf	50	75	LUPINS AND FIELD PEAS: Mixtures with Brodal or simazine may cause crop yellowing and separate applications are recommended. CHICKPEAS, FABO BEANS, LENTILS AND VETCH, LINOLA, LINSEED: Broadleaf herbicides should NOT be added to Wellfarm Haloxyfop 520 EC Herbicide. Apply Wellfarm Haloxyfop 520 EC Herbicide and broadleaf herbicides at least 1 week apart. LUCERNE, CLOVER OR MEDIC PASTURES: If grazed or cut for hay immediately prior to treatment delay application until all grasses have fully expanded leaves. Use 75 mL + spraying oil or 100 mL + wetter/ha. (See GENERAL INSTRUCTIONS, Spraying Oils/wetters section). If silver grass (Vulpia spp.) is present in pasture, simazine should be tank mixed with the higher rate of Wellfarm Haloxyfop 520 EC Herbicide plus a non-ionic wetter.
	early tillering	75	100	

Table 1B. Winter Crop Growth Stage Application Windows

Crop	Crop Growth Stage
Lucerne, Medic and Clover Pastures or Seed Crops	Apply from 2 nd trifoliate leaf onwards. For Erodium spp. spraying, apply from cotyledon crop stage onwards.
Canola, Linola, and Linseed	Apply from 2 leaf to 8 leaf stage of crop growth. DO NOT apply after the commencement of stem elongation. This means that application must NOT occur after the 8 leaf stage, or if stem elongation commences before the 8 leaf stage, application must NOT occur after stem elongation has commenced.
Chickpeas, Faba beans, Field peas, Lentils, Lupins, Vetch	Apply from 2 nd leaf, 2 nd node or 2 nd branch to prior to flowering.

Table 2A. Lucerne, Medic And Clover Seed Crops And Pastures - See Table 1b For Crop Stages

Weeds Controlled	Weed Growth Stage	Rate (mL/ha) with Uptake ¹ Spraying Oil	Critical Comments
Prairie grass (Bromus catharticus)	Up to early tillering	100	¹ See GENERAL INSTRUCTIONS, Spraying oils/wetters section.
Musky or ferny leaf Storksbill: (Erodium moschatum) Common Crowsfoot or Common Storksbill (Erodium cicutarium)	Up to 6 leaf or 5cm diameter	50 – 75 ³	³ Use lower rate when growing conditions and crop or pastures competition are good and when weed populations are below 100 plants/m ² . Use the higher rate when weed population are above 100 plants/m ² or when crop or pasture competition is poor.
Long or shiny leaf Storksbill (E. botrys)	Up to 8 leaf or 5cm diameter	75 – 100	NOTE: Storksbill may NOT be controlled if simazine or Broadstrike* are tank-mixed with Wellfarm Haloxyfop 520 EC Herbicide. LUCERNE, CLOVER OR MEDIC PASTURES If grazed or cut for hay immediately prior to treatment delay application until all grasses have fully expanded leaves. Use 75mL + spraying oil or 100mL + wetter/ha. (See GENERAL INSTRUCTIONS, Spraying oils/wetter section). If Silver grass (Vulpia spp.) is presented in pasture, simazine should be tank mixed with the higher rate of Wellfarm Haloxyfop 520 EC Herbicide plus a no-ionic wetter.

Table 2B. Lucerne, Medic And Clover Seed Crops Only (Not To Be Used For Stockfeed) - See Table 1B For Crop Stages

Weeds Controlled	Weed Growth Stage	Rate (mL/ha) with Uptake ¹ Spraying Oil	Critical Comments
Couch grass (suppression), Rhodes grass (control)	Tillering seedlings	150 + 150 ⁴	⁴ For best suppression of couch or control of Rhodes grass, make 2 applications of Wellfarm Haloxyfop 520 EC Herbicide 2-4 weeks apart. Time second application to coincide with tillering stage of weeds and just after irrigation or significant rain. Only treat actively growing weeds which are NOT moisture stressed. Use these rates for control of couch and Rhodes grass.
Couch grass (control), Rhodes grass (control)	Established stands	400 – 800	

Table 3A. Summer Crops – Cotton, Cowpea, Lucerne, Mung bean, Navy Beans, Peanuts, Soybeans, Sunflowers

Weeds Controlled	Weed Growth Stage	Rate (mL/ha) with Uptake ¹ Spraying Oil	Critical Comments
Australian millet	2 leaf to tillering up to 15cm	150	¹ See GENERAL INSTRUCTIONS, Spraying oils/wetters section. NAVY BEAN, PEANUTS, SOYBEANS For broadleaf weed control, Wellfarm Haloxyfop 520 EC Herbicide at 150mL/ha plus wetter may be tank mixed with Blazer® (except on Navy beans) or Basagran. Tank mixtures may cause transient leaf spotting on the crop but DO NOT normally affect yield. DO NOT tank mix broadleaf herbicides with Wellfarm Haloxyfop 520 EC Herbicide if grasses have begun tillering or if the grasses are under moisture stress. DO NOT add Uptake Spraying Oil when mixing with Blazer or Basagran. DO NOT use Blazer or Basagran tank-mixes on Cowpeas.
Barnyard grass	2 to 5 leaf	100	
	tillering up to 15cm	150	
Crowsfoot grass Green Panic grass Johnson grass (Rhizome)	2 leaf to tillering up to 15cm	150	
Johnson grass (seedling)	2 to 5 leaf	100	
Liverseed grass (seedling) Mossman River grass	tillering up to 15cm	150	
Summer grass	2 leaf to tillering up to 15cm	150	
Volunteer cereals	2 to 4 leaf	100	
	tillering up to 15cm	150	

Table 3B. Summer Crop Growth Stage Application Windows

Crop	Crop Growth Stage
Lucerne	Apply from 2 nd trifoliate leaf onwards
Cowpea, Mung beans, Navy beans, Soybeans	Apply from 2 nd leaf to flowering
Peanuts	Apply from 5cm to pegging
Cotton	Apply from 2 nd leaf to before the onset of flowering
Sunflowers	Apply from 2 nd leaf to head initiation

Table 4. Annual And Perennial Grasses And Erodium Spp. In Orchard, Vine, And Plantation Crops, Forestry And Pyrethrum

Crop	Crop Growth Stage	Weeds Controlled	Weed Growth Stage	Rate (mL/ha) with Uptake ¹ Spraying Oil	Critical Comments
Orchard, Vine and plantation Crops including: Apples, Avocado, Banana, Blueberry, Citrus, Custard apple, Feijoa, Grapevines, Guava, Kiwifruit, Litchi (Lychee), Longan, Mango, Nashi, Nut trees, Passionfruit, Paw paw, Pear, Persimmon, Pineapple, Rambutan, Stone Fruit	All growth stages	Perennial Grasses: Couch, Rhodes grass, Slender Rats Tail grass	Established stands	400 - 800	¹ See GENERAL INSTRUCTIONS, Spraying oils/wetters section. Spray should be directed to the base of the tree or vine avoiding contact with fruit and foliage. Spot Spray: Use 25mL to 50mL/100L of water. Use higher rate on late tillering mature grasses. Annual Grasses: Where treated in association with perennial grasses, these annual grasses will be controlled.
		Buffel grass, Green panic, Johnson grass, Kikuyu, Paspalum spp., Setaria spp.	Vegetative to early tillering	200	
			Late tillering	400	
		Annual Grasses: Annual ryegrass, Barley grass, Barnyard grass, Brome grass, Crowsfoot grass, Lesser Canary grass, Liverseed grass, Mossman river grass, Paradoxa grass, Summer grass, Volunteer cereals, Wild Oats	2 leaf to tillering	200	
Forestry: Pinus radiata, Eucalyptus spp.					

Crop	Crop Growth Stage	Weeds Controlled	Weed Growth Stage	Rate (mL/ha) with Uptake ¹ Spraying Oil	Critical Comments
Orchard, Vine and plantation Crops including: Apples, Avocado, Banana, Blueberry, Citrus, Custard apple, Feijoa, Grapevines, Guava, Kiwifruit, Litchi (Lychee), Longan, Mango, Nashi, Nut trees, Passionfruit, Paw paw, Pear, Persimmon, Pineapple, Rambutan, Stone Fruit	All growth stages	Perennial Grasses: Couch, Rhodes grass, Slender Rats Tail grass	Established stands	400 - 800	¹ See GENERAL INSTRUCTIONS, Spraying oils/wetters section. Spray should be directed to the base of the tree or vine avoiding contact with frit and foliage. Spot Spray: Use 25mL to 50mL/100L of water. Use higher rate on late tillering mature grasses. Annual Grasses: Where treated in association with perennial grasses, these annual grasses will be controlled.
		Buffel grass, Green panic, Johnson grass, Kikuyu, Paspalum spp., Setaria spp.	Vegetative to early tillering	200	
			Late tillering	400	
		Annual Grasses: Annual ryegrass, Barley grass, Barnyard grass, Brome grass, Crowsfoot grass, Lesser Canary grass, Liverseed grass, Mossman river grass, Paradoxa grass, Summer grass, Volunteer cereals, Wild Oats	2 leaf to tillering	200	
Forestry: Pinus radiata, Eucalyptus spp.					
Forestry: Pinus pineaster		Annual Grasses as above	Vegetative to tillering	125 - 250	Forestry: For annual grasses apply lowest rate to newly emerged grasses, increasing the rate as they develop.
Pyrethrum		Barley grass, Brome grass, Rope Twitch, Barnyard grass, Erodium spp., Volunteer cereals	Vegetative to tillering	100 - 250	Pyrethrum TAS only: For Erodium spp. apply 75-100 mL/ha if the main weed is E. botrys. Use 50-75 mL/ha if either E. cicutarium or E. moschatum are the main weeds.

Table 5. Wellfarm Haloxyfop 520 EC Herbicide And Select Herbicide Tank Mixes – Canola, Chickpeas, Faba Beans, Field Peas, Lupins, Lentils

Weeds Controlled	Weed Growth Stage	Rate (mL/ha)		Critical Comments
		Wellfarm Haloxyfop 520 EC Herbicide	Select Herbicide	
FOP/DIM Susceptible Annual Ryegrass + Volunteer barley, Volunteer wheat, Brome grass, Wild Oats, Barley grass, Phalaris	2 to 4 leaf	25	150	¹ See GENERAL INSTRUCTIONS, Spraying oils/wetters section. Use Uptake Spraying Oil at 500mL/100L or Hasten at 1L/100L. Apply at the same crop growth stages as those in Table 1B Winter Crops.
	Early tillering	38	150	
FOP Resistant Annual Ryegrass + Volunteer barley, Volunteer wheat, Brome grass, Wild Oats, Barley grass, Phalaris	2 to 4 leaf	25	200	Lentils: Apply up to 7 node-early branching crop growth stage only. Lupins: NOT for QLD.
	Early tillering	38	250	

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

HARVESTING WITHHOLDING PERIODS:

NOT REQUIRED WHEN USED AS DIRECTED FOR:

CANOLA, CHICKPEAS, COTTON, COWPEA, FABA BEANS, FIELD PEAS, LENTILS, LINOLA, LINSEED, LUPINS, MUNG BEANS, NAVY BEANS, ORCHARD CROPS, PEANUTS, PLANTATION CROPS, SOYBEANS, SUNFLOWERS, VETCH OR VINE CROPS.

DO NOT HARVEST FOR:

MEDIC AND CLOVER SEED CROPS	7 DAYS AFTER APPLICATION.
------------------------------------	----------------------------------

STOCK FOOD WITHHOLDING PERIODS:

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR:

CANOLA, CHICKPEAS, COTTON, COWPEA, FABA BEANS, FIELD PEAS, LENTILS, LINOLA, LINSEED, LUPINS, MUNG BEANS, NAVY BEANS, PEANUTS, SOYBEANS, SUNFLOWERS AND VETCH	28 DAYS AFTER APPLICATION.
---	-----------------------------------

LUCERNE	21 DAYS AFTER APPLICATION.
----------------	-----------------------------------

MEDIC AND CLOVER PASTURE	7 DAYS AFTER APPLICATION.
---------------------------------	----------------------------------

COTTON GIN TRASH MUST NOT BE FED TO ANIMALS.

GENERAL INSTRUCTIONS

MIXING

- Add water to the spray tank to 10cm above the level of agitation and ensure the agitation device is working vigorously (There must be a minimum of 100L of water in the tank before any pesticide is added).
- If tank mixing, firstly, add any soluble liquid formulations (e.g. Lontrel Herbicide) and allow agitation for approximately 1 minute.
- Then add Wellfarm Haloxyfop 520 EC Herbicide at the point where the agitation is the strongest (DO NOT add Wellfarm Haloxyfop 520 EC Herbicide through a strainer or sieve). Allow further agitation for 1 minute.
- Half fill the spray tank.
- If using wettable powder or water dispersible granules, or other emulsifiable concentration formulations (e.g. Lorsban 750 WG or Le-Mat), these should be added after the Wellfarm Haloxyfop 520 EC Herbicide to the half full spray tank ensuring vigorous agitation.
- Finally add Uptake Spraying Oil or approved alternate spraying oil/wetter (see section on spraying oils/wetters) and continue filling the tank to the required volume maintaining agitation at all times.
- Only mix sufficient solution for immediate use. Wellfarm Haloxyfop 520 EC Herbicide and any other tank mixes should be applied immediately for best results.

Spraying Oils/Wetters

1. **Spraying Oils:** It is essential to add an adjuvant to Wellfarm Haloxyfop 520 EC Herbicide. Best results will be achieved with Uptake Spraying Oil at 0.5L/100L of spray solution. Alternatively, other oils plus a non-ionic wetter may also be used. When other crop spraying oils are used, mix at 1L/100L and add a non-ionic wetter (surfactant) at 200mL/100L of spray solution. Use of an oil is NOT always recommended. See CRITICAL COMMENTS for specific situation recommendations.
2. **Non-Ionic Wetters:** When Uptake or other oils are NOT used, a 100% concentrate non-ionic wetting agent such as BS-1000 at 200mL/100L must be used along with the higher rate of Wellfarm Haloxyfop 520 EC Herbicide as specified in the DIRECTIONS FOR USE TABLE.

Where Water Volumes Of Less Than 50L/Ha Are Used, DO NOT Use Less Than 250mL/Ha Of Uptake Or 500mL/Ha For Oils Other Than Uptake Or Less Than 100mL/Ha Of Wetter.

CANOLA, LUCERNE, MEDIC AND CLOVER PASTURES AND SEED CROPS

When tank mixing Wellfarm Haloxyfop 520 EC Herbicide with Lontrel herbicides (canola only) or Broadstrike* (lucerne, clover and medics), use Uptake Spraying Oil with the lower rates of Wellfarm Haloxyfop 520 EC Herbicide or a wetting agent with the higher rates of Wellfarm Haloxyfop 520 EC Herbicide unless otherwise specified. When mixing Wellfarm Haloxyfop 520 EC Herbicide with other broadleaf herbicides on these crops, DO NOT use an oil use a wetter instead.

FIELD PEAS AND CANOLA

The oil recommended is Uptake Spraying Oil. Hasten is also recommended for use with tank-mixtures of Wellfarm Haloxyfop 520 EC Herbicide and Select Herbicide.

For canola, Wellfarm Haloxyfop 520 EC Herbicide + Lontrel 750 SG + Uptake Spraying Oil are compatible and selective to canola. This tank-mixture is also compatible with atrazine or simazine and selective to triazine tolerant canola.

NAVY BEANS, PEANUTS AND SOYBEANS

When mixing with Blazer or Basagran DO NOT add spraying oil to these mixtures. DO NOT use these tank-mixtures on cowpea.

COMPATIBILITY

Ground Use Only: Wellfarm Haloxyfop 520 EC Herbicide Can Be Tank-Mixed With:

Insecticides:	Dimethoate, Lorsban 500 EC Insecticide, Lorsban 750 WG Insecticide, Omethoate
Herbicides:	Atrazine, Basagran, Blazer, Broadstrike* Herbicide, Lontrel Herbicide, Lontrel 750 SG, MCPA Ester (LVE) – (DO NOT exceed 700mL/ha of MCPA LVE), Oryzalin, Select Herbicide, Simazine, Starane 200 Herbicide
Fungicides:	Dithane DF, Dithane Rainshield Fungicide
Trace Elements:	Magnesium Sulphate, Zinc Sulphate

Wellfarm Haloxyfop 520 EC Herbicide Is NOT Compatible With 2,4-D Or MCPA As Sodium Or Amine Salts.

* Other Registered Trademarks

Aerial Use Only

No product other than a recommended crop oil or wetter should be mixed with Wellfarm Haloxyfop 520 EC Herbicide when applied by air except for the addition of Lontrel Herbicide for use in forestry and Lontrel 750 SG for use in canola only.

APPLICATION

Apply Wellfarm Haloxyfop 520 EC Herbicide in sufficient water to obtain good coverage. It should be applied by an accurately calibrated ground rig or aircraft delivering droplets with a VMD of 200-300 microns.

The Following Spray Volumes Are Recommended:

Ground Application: 50-150 L/ha

Aerial Application: 30 L/ha minimum

Use Higher Water Volumes In Orchards And In Dense Crops Where The Weeds May Be Shielded By The Crop Canopy.

CLEANING SPRAY EQUIPMENT

Before Using Wellfarm Haloxyfop 520 EC Herbicide

If broadleaf herbicides, particularly sulfonylureas, have been used in the spray equipment at any time prior to the use of Wellfarm Haloxyfop 520 EC Herbicide, particular care should be taken to follow the directions on the relevant broadleaf herbicide label for equipment cleaning, or damage to susceptible crops may occur.

After Using Wellfarm Haloxyfop 520 EC Herbicide

Empty the tank completely and drain the whole system. Thoroughly wash inside the tank using a pressure hose, drain the tank and clean any filters in the tank, pump, line and nozzles.

To Rinse: After cleaning the tank as above, quarter fill the tank with clean water and circulate through the pump, lines, hoses and nozzles. Drain and repeat the rinsing procedure twice.

To Decontaminate: Before spraying cereals, maize, sorghum or other sensitive crops, wash the tank and rinse the system as above. Then quarter fill the tank and add an alkali detergent (e.g. SURF, Cold Water SURF Concentrate, DynamoMatic Concentrate, OMO or Drive) at 500mL/100L of water or the powder equivalent at 500g/100L of water, and circulate throughout the system for at least 15 minutes. Drain the whole system. Remove filters and nozzles and clean them separately. Finally flush the system with clean water and allow to drain. Chlorine based cleaners are NOT recommended. Rinse water should be discharged onto a designated disposal area, or if this is unavailable, onto unused land away from desirable plants and water sources.

RESISTANT WEEDS WARNING

GROUP	A	HERBICIDE
--------------	----------	------------------

Wellfarm Haloxyfop 520 EC Herbicide is a member of the aryloxyphenoxy propionate group of herbicides. Wellfarm Haloxyfop 520 EC Herbicide has the acetyl CoA carboxylase inhibitor mode of action. For weed resistance management, Wellfarm Haloxyfop 520 EC Herbicide is a Group A herbicide.

Some naturally occurring weed biotypes resistant to Wellfarm Haloxyfop 520 EC Herbicide and other Group A herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if Wellfarm Haloxyfop 520 EC Herbicide and other Group A herbicides are used repeatedly. These resistant weeds will NOT be controlled by Wellfarm Haloxyfop 520 EC Herbicide or any other Group A herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Wellfarm Pty Ltd accepts NO liability for any losses that may result from the failure of Wellfarm Haloxyfop 520 EC Herbicide to control resistant weeds.

Wellfarm Haloxyfop 520 EC Herbicide may be subject to specific resistance management strategies. For further information consult your local farm chemical supplier, consultant, local Department of Agriculture or Primary Industries.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Wellfarm Haloxyfop 520 EC Herbicide damages cereals and grasses.

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

Cereal crops or grasses planted within 12 weeks of application may be damaged by the residual effects of Wellfarm Haloxyfop 520 EC Herbicide, particularly on light and red soils.

PROTECTION OF LIVESTOCK

DO NOT graze or cut treated crops for stock food except as specified under withholding periods.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

TOXIC TO FISH.

DO NOT contaminate dams, streams, drains, rivers or waterways with the chemicals or used containers.

STORAGE AND DISPOSAL

KEEP OUT OF REACH OF CHILDREN

Store in the closed, original container in a dry, cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT store near feedstuffs, fertilisers or seeds. Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty containers for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty containers 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory Government Regulations. DO NOT burn empty containers or product.

SMALL SPILL MANAGEMENT

Wear protective equipment (see SAFETY DIRECTIONS). Apply absorbent material such as earth, sand, cat litter or clay granules to the spill. When absorption is complete, sweep up material and contain in a refuse vessel for disposal (see STORAGE AND DISPOSAL section). If necessary wash the spill area with an alkali detergent and water and absorb this wash liquid for disposal as directed above.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening container, preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and washable hat, elbow-length PVC gloves and face shield or goggles. After each day's use, wash gloves and face shield or goggles and contaminated clothing. Wash hands after use.

FIRST AID

If poisoning occurs contact a Doctor or phone the Australian Poisons Information Centre 13 11 26. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a Doctor.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS) that can be obtained from the supplier.

CONDITIONS OF SALE

The use of this product is beyond the control of the manufacturer. No warranty expressed or implied is given by Wellfarm Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Wellfarm Pty Ltd accepts NO responsibility for any consequences whatsoever resulting from the use of this product.

MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY

Wellfarm Pty Ltd has no control over storage, handling and manner of use of this product. Where the material is not stored, handled or used correctly and in accordance with the directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory

warranties) will apply. Wellfarm Pty Ltd accepts NO liability for any losses or damages arising from incorrectly storage, handling or use.

IN AN EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE

APVMA Approval No 70164/0714

Wellfarm Pty Ltd
ACN 158 518 551
22 Calypso Crescent
Point Cook Vic 3030
www.wellfarm.com.au