

**CAUTION** KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



# Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC

## Herbicide

**ACTIVE CONSTITUENT: 25g/L DIFLUFENICAN**  
**250g/L MCPA PRESENT AS THE ETHYL HEXYL ESTER**  
**SOLVENT: 325g/L HYDROCARBON LIQUID**  
**150g/L N-METHYL-2-PYRROLIDONE**

**GROUP FI HERBICIDE**

For the control of certain broadleaf weeds in winter cereals and clover  
as specified in the Directions For Use Table

**IMPORTANT: READ THE ATTACHED LEAFLET  
BEFORE OPENING OR USING THIS PRODUCT**

CONTENTS

**20L**

### STORAGE AND DISPOSAL KEEP OUT OF REACH OF CHILDREN

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight.

#### For Only (5L, 20L)

Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty containers for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty containers 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory Government Regulations. DO NOT burn empty containers or product.

#### For Refillable Containers

If tamper evident seals are broken prior to initial use then the integrity of the contents CANNOT be assured. Empty product as required into application equipment. DO NOT attempted to breach the valve system or filling point, or contaminate the container with water or other products. Ensure that equipment used in transfer of the product is disconnected, triple rinsed and drained after each use. When the container is empty, close all caps and valves and return the container to the point of purchase.

### SAFETY DIRECTIONS

Harmful if swallowed. Will damage eyes. Will irritate the skin. Avoid contact with eyes and skin. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and face shield or goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield or goggles and contaminated clothing.



Wellfarm Pty Ltd  
ACN 158 518 551  
ABN 32 158 518 551

22 Calypso Crescent  
Point Cook Vic 3030  
www.wellfarm.com.au

### FIRST AID

If poisoning occurs contact a Doctor or phone the Australian Poisons Information Centre 13 11 26. If swallowed, DO NOT induce vomiting. Give a glass of water. If in eyes, wash out immediately with water.

### MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS) that can be obtained from the supplier.

### CONDITIONS OF SALE

The use of this product is beyond the control of the manufacturer. No warranty expressed or implied is given by Wellfarm Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Wellfarm Pty Ltd accepts NO responsibility for any consequences whatsoever resulting from the use of this product.

### MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY

Wellfarm Pty Ltd has no control over storage, handling and manner of use of this product. Where the material is not stored, handled or used correctly and in accordance with the directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Wellfarm Pty Ltd accepts NO liability for any losses or damages arising from incorrectly storage, handling or use.

**APVMA Approval No 70185/62692**

**IN AN EMERGENCY  
DIAL 000 POLICE  
OR FIRE BRIGADE**



Batch No

DOM

# CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

## Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide

ACTIVE CONSTITUENT: 25g/L DIFLUFENICAN  
250g/L MCPA PRESENT AS THE ETHYL HEXYL ESTER  
SOLVENT: 325g/L HYDROCARBON LIQUID  
150g/L N-METHYL-2-PYRROLIDONE

GROUP	<b>FI</b>	HERBICIDE
-------	-----------	-----------

For the control of certain broadleaf weeds in winter cereals and  
clover as specified in the Directions For Use Table

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE  
OPENING OR USING THIS PRODUCT**

**CONTENTS: 20L**  
(5L, 100L, 110L, 1000L)

APVMA Approval No 70185/62692

Wellfarm Pty Ltd  
ACN 158 518 551  
22 Calypso Crescent  
Point Cook Vic 3030  
[www.wellfarm.com.au](http://www.wellfarm.com.au)



**WEED LIST**

<b>WEED</b>	<b>Common Name</b>	<b>Scientific Name</b>
	Canola (Rape)	Brassica Napus
	Capeweed	Arctotheca Calendula
	Charlock	Sinapis Arvensis
	Chickweed	Stellaria Media
	Common Sowthistle (Milk Thistle)	Sonchus Oleraceus
	Corn Gromwell	Buglossoides Arvense
	Cowvine	Ipomoea Lonchophylla
	Crassula	Crassula Spp.
	Deadnettle	Lamium Amplexicaule
	Dense-Flower Fumitory	Fumaria Densiflora
	Dock	Rumex Spp.
	Doublegee (Spiny Emex)	Emex Australis
	Fat Hen	Chenopodium Album
	Fireweed	Senecio Spp.
	Fumitory	Fumitory Spp.
	Hedge Mustard	Sisymbrium Officinale
	Hexham Scent (King Island Melilot)	Melilotus Indicus
	Horehound	Marrubium Vulgare
	Hyssop Loosestrife	Lythrum Hyssopifolia
	Iceplant	Mesembryanthemum Spp.
	Indian Hedge Mustard	Sisymbrium Orientale
	London Rocket	Sisymbrium Irio
	Long Storksbill	Erodium Botrys
	Marshmallow	Malva Parviflora
	Mouse-Eared Chickweed	Cerastium Glomeratum
	Night-Scented Stock	Matthiola Longipetala
	Paterson's Curse	Echium Plantagineum
	Peppercress	Lepidium Spp.
	Prickly Lettuce	Lactuca Serriola
	Purple Goosefoot	Scleroblitum Atriplicinum
	Rough Poppy	Papaver Hybridum
	Saffron Thistle	Carthamus Lanatus
	Scarlet Pimpernel	Anagallis Arvensis
	Shepherd's Purse	Capsella Bursa-Pastoris
	Skeleton Weed	Chondrilla Juncea
	Sorrel	Rumex Spp.
	Stemless Thistle	Onopordum Acaulon
	Toad Rush	Juncus Bufonius
	Tree Hogweed	Polygonum Patulum
	Turnip Weed	Rapistrum Rugosum
	Varigated Thistle	Silybum Marianum
	Vetch (Tares)	Vicia Sativa
	Volunteer Lupins	Lupinus Spp.
	Ward's Weed	Carrichtera Annua
	Wild Radish	Raphanus Raphanistrum
	Wild Turnip	Brassica Tournefortii
	Wireweed (Hogweed)	Polygonum Aviculare

## DIRECTIONS FOR USE RESTRAINTS

DO NOT apply to crops or weeds which are stressed due to dry or excessively moist conditions.

DO NOT apply to crops under stress due to disease or insect damage.

DO NOT apply to frost affected crops or if frosts are imminent.

DO NOT apply when heavy rain is expected within 4 hours.

CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE /ha	CRITICAL COMMENTS
CEREALS Wheat, Barley, Oats, Triticale, Cereal Rye (including cereals undersown with clover)  PASTURE Newly Sown and Established Clover-Based Pasture, Clover for Hay and Seed Production	Wild Radish	Up to the 2 leaf stage and not more than 60mm in diameter	WA only	250mL	CROP STAGE Cereals Up to 750mL (3 leaf to late tillering stage - Z13 to 30) Over 750mL (5 leaf to late tillering stage - Z15 to 30) Optimum results are achieved when sprayed at 3-5 leaf crop stage (generally 4-8 weeks post sowing).  WA only: DO NOT apply to Barley or Kulin Wheat before the 5 leaf stage (Z15).  Warning: Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide may cause transient crop yellowing of cereals. Some varieties of oats have NOT been tested. Refer to 'Crop Tolerance' section of General Instructions.  Application is recommended prior to the eighth trifoliate leaf stage, however, applications prior to the third leaf stage may result in crop damage especially under stressed conditions and in sandy soils. DO NOT apply to Annual Medics or Lucerne.  Warning: Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide may cause transient crop yellowing of clover, and may affect growth and seed set of some varieties of clover. Refer to 'Crop Tolerance' section of General Instructions.  WEED STAGE Apply when weeds are actively growing. In most situations the rate specified for each weed size will give satisfactory control. Under certain conditions such as: • high crop and weed density. • late season germinations. • abnormal weed growth (including early flowering), high rate of product (up to the maximum rate of application specified for that weed) may be required.
		Up to the 4 leaf stage and not more than 120mm in diameter	All States	500mL	
		Up to the 6 leaf stage and not more than 150mm in diameter		750mL	
		Up to the 8 leaf stage and not more than 180mm in diameter		1.0L	
	Charlock, Hedge Mustard, Indian Hedge Mustard, Shepherd's Purse, Turnip Weed, Wild Turnip	500mL			
		Up to the 4 leaf stage and not more than 120mm in diameter		750mL	
		Up to the 6 leaf stage and not more than 150mm in diameter		1.0L	
	London Rocket	Up to the 5 leaf stage and not more than 120mm in diameter	QLD only	750mL	
	Ward's Weed		SA only		
	Capeweed	Up to the 2 leaf stage and not more than 60mm in diameter	All States	500mL	
		Up to the 4 leaf stage and not		1.0L	

CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE /ha	CRITICAL COMMENTS
		more than 120mm in diameter			Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide will NOT effectively control: <ul style="list-style-type: none"><li>• regrowth of suppressed weeds;</li><li>• transplanted weeds;</li><li>• regrowth from rhizomes or roots;</li><li>• weeds growing under stress from previous herbicide applications.</li></ul> GRAZING Efficacy on larger weeds will be improved by grazing with normal levels of stock after 7 day withholding period. Refer to 'Protection of Livestock' for grazing precautions.  APPLICATION Activity of this product will be reduced if weeds are stressed. Optimum results will be obtained if good soil moisture exists at and after application. Where crop or weed density is high, water volume should be increased.  WILD RADISH Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide will provide residual control of Wild Radish for up to 4 weeks after application. Effective residual activity of this product may be reduced where: <ul style="list-style-type: none"><li>• rates lower than 1.0L/ha are used;</li><li>• dry conditions prevail;</li><li>• poor coverage of the soil surface is achieved;</li><li>• crop is planted in non-wetting sand;</li><li>• soils contain a high content of organic matter.</li></ul> Optimum results will be obtained if good soil moisture exists at and after application.
	Crassula	Up to the 2 leaf stage		500mL	
		Up to the 4 leaf stage		750mL	
	Prickly Lettuce	Up to the 2 leaf stage		500mL	
		Up to the 4 leaf stage		750mL	
		Up to the 6 leaf stage		1.0L	
	Dense-Flower Fumitory	Up to the 2 leaf stage		750mL	
	Corn Gromwell, Saffron Thistle, Toad Rush			1.0L	
	Deadnettle		NSW, VIC, SA only	1.0L	
	Sorrel	Up to 2 leaf stage	VIC only	1.0L	
	Canola (Rape)	Up to 4 leaf stage	All States	500mL	
	Purple Goosefoot	Up to 6 leaf stage	QLD only	500mL	
	Turnip Weed Wild Turnip	Cotyledon to 2 leaf	NSW only (West of Newell Hwy), SA only (Eyre Peninsula North of the Line between Venus Bay and Cowell)	350mL	
CEREALS Wheat, Barley, Oats, Triticale, Cereal Rye	Fumitory	2 – 6 leaf stage	All States	500 + 200mL terbutryn (500g/L)	

# SUPPRESSION OF THE FOLLOWING WEEDS

CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE /ha	CRITICAL COMMENTS
CEREALS Wheat, Barley, Oats, Triticale, Cereal Rye (including cereals undersown with clover)  PASTURE Newly Sown and Established Clover-Based Pasture, Clover for Hay and Seed Production	Saffron Thistle	Up to 6 leaf stage	All States	1.0L	Refer to critical comments in the previous section.
	Chickweed, Fireweed, Hexham Scent (King Island Melilot), Iceplant, Mouse-Eared Chickweed, Night-Scented Stock, Paterson's Curse, Peppergrass, Skeleton Weed, Long Storksbill, Volunteer Lupins.	Up to 4 leaf stage			
	Wireweed (Hogweed)			750mL	
	Common Sowthistle (Milk Thistle), Cowvine, Dock, Doublegee, (Spiny Emex), Fat Hen, Horehound, Hyssop, Loosestrife, Marshmallow, Rough Poppy, Scarlet Pimpernel, Stemless Thistle, Tree Hogweed, Variegated Thistle, Vetch (Tares).	Up to 2 leaf stage		1.0L	

CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE /ha	CRITICAL COMMENTS
CEREALS Wheat, Barley, Oats, Triticale, Cereal Rye	Wild Radish	Up to the 4 leaf stage and not more than 120mm in diameter	All States	350mL plus 200mL L.V.E. MCPA (500g/L)	Refer also to all Critical Comments relating to weed stage, grazing, application and wild radish above. *Reduced efficacy (suppression only) may be achieved on wild radish larger than 8 leaf or greater than 180mm in diameter. DO NOT use this tank-mix if cereals

		Up to the 6 leaf stage and not more than 150mm in diameter		500mL plus 200mL L.V.E. MCPA (500g/L)	are undersown with lucerne or annual medics. Crop Stage Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide 350mL + L.V.E. MCPA 200mL: Apply from 3 leaf to fully tillered (Zadoks Z13 to Z30).
		Up to the 8 leaf stage and not more than 180mm in diameter*		500mL plus 400mL L.V.E. MCPA (500g/L)	Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide 500mL + L.V.E. MCPA 200mL: Apply from 3 leaf to fully tillered (Zadoks Z13 to Z30). Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide 500mL + L.V.E. MCPA 400mL: Apply from 5 leaf stage to fully tillered (Zadoks Z15 to Z30). Optimum results are achieved when sprayed at 3-5 leaf crop stage (generally 4-8 weeks post-sowing). WA only: DO NOT apply to Barley or Kulin Wheat before the 5 leaf stage (Z15). Warning: Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide may cause transient crop yellowing of cereals. Some varieties of oats have NOT been tested. Refer to 'Crop Tolerance' section of General Instructions. Observe instructions also on L.V.E. MCPA product label.

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL  
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIODS:**

<b>CROP HARVEST</b>	<b>NOT REQUIRED WHEN USED AS DIRECTED.</b>
<b>ALL CROPS</b>	<b>DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.</b>

**GENERAL INSTRUCTIONS**

## RESISTANT WEEDS WARNING

# GROUP F I HERBICIDE

Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide is a member of the phenoxy and nicotinanalide groups of herbicides and acts by inhibiting carotenoid biosynthesis and disrupting plant cell growth. For weed resistance management, Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide is both a Group F and a Group I herbicide.

Some naturally occurring weed biotypes resistant to Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide and other Group F and Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide and other Group F and Group I herbicides are used repeatedly. These resistant weeds will NOT be controlled by Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide or any other Group F and Group I herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Wellfarm Pty Ltd accepts NO liability for any losses that may result from the failure of Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide to control resistant weeds.

Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide may be subject to specific resistance management strategies. For further information consult your local farm chemical supplier, consultant, local Department of Agriculture or Primary Industries.

## TOLERANCE

Some pre-emergence herbicides, such as Atrazine, can cause stress to certain crops resulting in an increase in crop damage when using this product. Sub-clover is particularly sensitive.

## CEREALS

After application some transient crop yellowing may occur. This usually appears as yellow or white banding on leaves. Provided the crop is NOT under stress from pre-emergent herbicide, root disease, insect damage, frost, dry or excessively moist conditions, the development of the crop and subsequent growth will be unaffected.

## WARNING (Oats)

The tolerance of varieties Esk and Nile (the two main varieties grown in TAS) to Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide has NOT been tested. Test a small area of crop before using Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide over large areas. Consult your farm chemical supplier, consultant, local Department of Agriculture or Primary Industries, or local Wellfarm Pty Ltd representative for advice on specific varieties.

## PASTURE

The tolerance of clover varieties to Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide can vary with rate of application, soil type, crop health, stage of growth and degree of moisture and temperature stress.

## WARNING

Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide may result in transient crop yellowing and suppression of growth with a resultant initial reduction in dry matter, particularly at rates in excess of 500mL/ha and in areas of double spray. For this reason application is recommended prior to the 8<sup>th</sup> trifoliolate leaf stage. However, at the lower rates (500mL/ha and less) and under normal growing conditions, subsequent growth and seed yield should NOT be affected.

Under normal growing conditions, the following varieties have shown acceptable levels of foliage tolerance to Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide applied at 500mL/ha:

**Arrow Leaf:** Zulu

**Balansa:** Paradana

**Berseem:** Sacromonte

**Persian:** Kyambro, Lupers, Maral

**White:** Haifa

**Subterranean Clover:** Daliak, Dalkeith, Denmark, Esperance, Geraldton, Goulburn, Karridale, Larissa, Leura, Mt. Barker, Nungarin, Rosedale, Seaton Park, Trikkala and Woogenellup. The effects of



Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide on clover seed yield have been tested on the following varieties. Under normal growing conditions they show levels of tolerance to Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide applied at 500mL/ha.

**Subterranean Clover:** Esperance, Goulburn, Larissa, Seaton Park and Trikkala.

#### **WARNING**

Rose and Strawberry clover have shown increased sensitivity to Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide. Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide may affect the seed of yield of subterranean variety Woogenellup. Some pasture grasses, including Phalaris and Cocksfoot, may show some initial reduction in vegetative growth after application of Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide. Care should be exercised if sensitive clover varieties or grasses are included in the pasture sward. Varieties NOT listed should be tested before using Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide over large areas. Consult your farm chemical supplier, consultant, local Department of Agriculture or Primary Industries, or local Wellfarm Pty Ltd representative for advice on specific varieties.

#### **SUBSEQUENT CROPS**

To reduce effect on susceptible crops (e.g. canola), ensure thorough cultivation of soil prior to the sowing of these crops.

#### **MIXING**

To ensure even mixing, half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly then add the remainder of the water. Agitate again before spraying commences. Reseal part-used product container immediately after use. Spray mixtures containing Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide should NOT be left to stand overnight. Prolonged periods of exposure to cold temperature could result in settling out of the product in the mixture.

#### **WARNING**

The rubber components present in some spraying units may be affected by exposure to the solvents in Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide and some other agricultural products. To reduce the risk it is recommended that the spray unit be thoroughly washed with a boom cleaner and fresh water after use. Contact the spray unit manufacturer to determine the suitability of the rubber components for use with agricultural products.

#### **APPLICATION**

##### **Boom Sprayer**

A minimum of 50L of water per hectare should be used, however, for optimum results water rates of 70-100L/ha are recommended. Increase the water volume if weed infestation is heavy or crop cover is dense. Complete coverage of weeds is essential.

#### **SELECTIVE HERBICIDE**

##### **Aircraft (NSW, VIC, SA only)**

Apply in a minimum of 30L water per hectare. Effective control will only be achieved where good coverage of leaf surface is achieved.

#### **COMPATIBILITY**

The following products are physically compatible with Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide as a two-way mixture in the spray tank but should only be used for the crops specified:

<b>Crop</b>	<b>Wellfarm Diflufenican 25 + MCPA Iso- Octyl Ester 250 EC Herbicide</b>	<b>Compatible Product</b>
Wheat, Triticale and Cereal Rye only	Up to 750mL/ha	Hoegrass (also Barley), Puma Progress (Wild Oats only), Tristar Advance (1.5L rate only), Wildcat (Wild Oats only)
Cereals (including undersown)	All rates	Chlorpyrifos (500g/L), Dimethoate, Endosulfan

Cereals (not undersown)	Up to 500mL/ha	Metsulfuron, Chlorsulfuron, L.V.E. MCPA, Triasulfuron
	All rates	Bromoxynil (200g/L), Amine 625, Clopyralid, Tordon 50-D, Dicamba (900g/kg) (up to 115g only), Eclipse
Wheat, Barley, Triticale, and Cereal Rye only (not undersown)		Tralkoxydim
Wheat only (not undersown)		Clodinafop
Clover	Up to 750mL/ha	Targa, Fusilade
Sub Clover		Simazine liquid, Simazine liquid + Nuquat mixture
	Up to 1.0L/ha	2,4-DB Amine (500g/L)

When mixing with other herbicides, crop yellowing may be enhanced. When mixing with Hoegrass, Wildcat, Puma Progress or Tristar Advance some reduction in the efficacy and speed of action of these products may occur. If the crop is stressed, the application of the herbicide tank-mixtures may cause yield reduction. When mixing with dicamba dry a temporary wilting may be evident in some crops after application. Growers should seek advice before spraying recently released cereal varieties. Use the recommended rates for both herbicides in the tank-mixture as well as the surfactant recommendation of the grass herbicide. If another herbicide is applied as a tank mix, observe the plantback restrictions on that label. DO NOT add surfactant when mixing Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide and Metsulfuron.

**Simazine:** Refer to the simazine label for correct application rates, especially with regard to soil types.

**This Product May Be Mixed In The Spray Tank With One Of The Following Insecticides**

**According To The Directions For Use On The Product:** Hallmark 50EC, Dominex 100EC, Karate, Decis Forte EC, and Talstar.

## **WARNING**

DO NOT use crop oils with Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide or Wellfarm Diflufenican 25 + MCPA Iso-Octyl Ester 250 EC Herbicide mixtures with other products in cereals. As formulations of other manufacturer's products are beyond the control of Wellfarm Pty Ltd, all mixtures should be tested prior to mixing commercial quantities.

## **PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

DO NOT apply under weather conditions, or from spraying equipment, that may cause drift onto nearby Susceptible Plants/Crops such as Cotton, Tobacco, Tomatoes, Vines, Lupins, Fruit Trees and Ornamentals.

## **PROTECTION OF LIVESTOCK**

### **Grazing Precautions**

Sprayed weeds may become more palatable to stock and a higher intake of some weeds may result in stock poisoning and death from causes such as nitrate poisoning. Care should be taken especially where capeweed, Paterson's curse and variegated thistles predominate in the pasture. Avoid grazing with young or breeding stock. DO NOT graze horses or pigs on Paterson's curse. If in doubt, contact your nearest Department of Agriculture.

## **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

**DANGEROUS TO FISH.**

DO NOT contaminate dams, streams, drains, rivers or waterways with the chemicals or used containers.

## **STORAGE AND DISPOSAL**

### **KEEP OUT OF REACH OF CHILDREN**

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight.

**For Only (5L, 20L)**

Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty containers for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty containers 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory Government Regulations. DO NOT burn empty containers or product.

**For Refillable Containers**

If tamper evident seals are broken prior to initial use then the integrity of the contents CANNOT be assured. Empty product as required into application equipment. DO NOT attempted to breach the valve system or filling point, or contaminate the container with water or other products. Ensure that equipment used in transfer of the product is disconnected, triple rinsed and drained after each use. When the container is empty, close all caps and valves and return the container to the point of purchase.

**SAFETY DIRECTIONS**

Harmful if swallowed. Will damage eyes. Will irritate the skin. Avoid contact with eyes and skin. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and face shield or goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield or goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs contact a Doctor or phone the Australian Poisons Information Centre 13 11 26. If swallowed, DO NOT induce vomiting. Give a glass of water. If in eyes, wash out immediately with water.

**MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet (MSDS) that can be obtained from the supplier.

**CONDITIONS OF SALE**

The use of this product is beyond the control of the manufacturer. No warranty expressed or implied is given by Wellfarm Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Wellfarm Pty Ltd accepts NO responsibility for any consequences whatsoever resulting from the use of this product.

**MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY**

Wellfarm Pty Ltd has no control over storage, handling and manner of use of this product. Where the material is not stored, handled or used correctly and in accordance with the directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Wellfarm Pty Ltd accepts NO liability for any losses or damages arising from incorrectly storage, handling or use.

**IN AN EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE**

APVMA Approval No 70185/62692

Wellfarm Pty Ltd  
ACN 158 518 551  
22 Calypso Crescent  
Point Cook Vic 3030  
[www.wellfarm.com.au](http://www.wellfarm.com.au)