

POISON KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Bromoxynil 200 EC

Herbicide

**ACTIVE CONSTITUENT: 200g/L BROMOXYNIL
PRESENT AS THE N-OCTANOYL ESTER
SOLVENT: 616g/L HYDROCARBON LIQUID**

GROUP C HERBICIDE

For the control of weeds in wheat, oats, barley, cereal rye and triticale (alone or undersown), linseed, clover and lucerne pastures, turf, fallow, non-crop, roadsides and right-of-way as specified in the Directions For Use Table

**IMPORTANT: READ THE ATTACHED LEAFLET
BEFORE OPENING OR USING THIS PRODUCT**

CONTENTS
20L

GENERAL INSTRUCTIONS

Before opening, carefully read Directions for Use, Precautionary and Protection Statements, Storage and Disposal, Safety Directions and First Aid instructions. Use only at recommended growth stages of crops and weeds. DO NOT use on crops other than recommended.

STORAGE AND DISPOSAL KEEP OUT OF REACH OF CHILDREN For Only (5L, 20L, 200L)

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty containers for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty containers 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory Government Regulations. DO NOT burn empty containers or product.

For Refillable Containers (1000L)

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Product is poisonous if inhaled or swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing spray, wear elbow-length PVC gloves and face shield. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield.

FIRST AID

If poisoning occurs contact a Doctor or phone the Australian Poisons Information Centre 13 11 26. If swallowed, DO NOT induce vomiting. Give a glass of water.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS) that can be obtained from the supplier.

CONDITIONS OF SALE

The use of this product is beyond the control of the manufacturer. No warranty expressed or implied is given by Wellfarm Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Wellfarm Pty Ltd accepts NO responsibility for any consequences whatsoever resulting from the use of this product.

MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY

Wellfarm Pty Ltd has no control over storage, handling and manner of use of this product. Where the material is not stored, handled or used correctly and in accordance with the directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Wellfarm Pty Ltd accepts NO liability for any losses or damages arising from incorrectly storage, handling or use.

APVMA Approval No 70239/62804

**IN AN EMERGENCY
DIAL 000 POLICE
OR FIRE BRIGADE**



Wellfarm Pty Ltd
ACN 158 518 551
ABN 32 158 518 551
22 Calypso Crescent
Point Cook Vic 3030
www.wellfarm.com.au



Batch No

DOM

POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

Wellfarm Bromoxynil 200 EC Herbicide

**ACTIVE CONSTITUENT: 200g/L BROMOXYNIL
PRESENT AS THE N-OCTANOYL ESTER
SOLVENT: 616g/L HYDROCARBON LIQUID**

GROUP	C	HERBICIDE
--------------	----------	------------------

For the control of weeds in wheat, oats, barley, cereal rye and triticale (alone or undersown), linseed, clover and lucerne pastures, turf, fallow, non-crop, roadsides and right-of-way as specified in the Directions For Use Table

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE
OPENING OR USING THIS PRODUCT**

**CONTENTS: 20L
(5L, 200L, 1000L)**

APVMA Approval No 70239/62804

Wellfarm Pty Ltd
ACN 158 518 551
22 Calypso Crescent
Point Cook Vic 3030
www.wellfarm.com.au



DIRECTIONS FOR USE

RESTRAINTS

DO NOT apply by mister machines.

DO NOT apply if crops or weeds are stressed due to excessively dry or moist conditions.

DO NOT apply to diseased or frost-affected crops or if frosts are imminent.

DO NOT apply if rain is expected within 3 hours.

DO NOT apply to lucerne crops or clover-based pastures or turf in temperatures above 20°C or when temperatures above 20°C may follow for some days after application.

Situation & Crop	Weeds Controlled	Weed Stage	State	Rate	Critical Comments
Wheat, Barley, Cereal Rye, Oats and Triticale: including those undersown with Clover, Lucerne or Medic (VIC only); Linseed; Clover or Lucerne Pastures; Lotus Seed or Forage Crops (QLD only)	Amsinckia (Amsinckia spp.), Capeweed (Arctotheca calendula) Black Bindweed (Climbing Buckwheat) (Fallopia convolvulus) Chamomile (Matricaria matricarioides), Common Peppergrass (Lepidium africanum), Corn Gromwell (Ironweed, Sheepweed), (Buglossoides arvensis) Fat Hen (Chenopodium album), Field Madder (Sherardia arvensis), Lesser Swinecress (Coronopus didymus), Mountain Sorrel (Oxalis acetosella) Purple Calandrinia (Calandrinia menziesii), Saffron Thistle (Carthamus lanatus), Shepherd's Purse (Capsella bursa-pastoris), Three-horned bedstraw (Galium trichomutum) Tree Hogweed (Polygonum patulum), Variegated Thistle (Silybum marianum)	Up to the 4 leaf stage or when plants are no more than 35 mm in diameter	All states	1.4 L/ha	<p>CROP STAGE: Wheat, Barley, Cereal, Rye, Oats and Triticale: 3 leaf to fully tillered (Zadoks Scale Z13-30). Linseed: 50 to 150 mm high. Lucerne and Clover: After seedlings have one trifoliate leaf or before there is 150mm regrowth in established stands. Lotus: Apply post-emergence at least 4 weeks after sowing (crop seedlings with a min. of 2-6 true leaves).</p> <p>APPLICATION: Refer to General Instructions. Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application. Good coverage of the spray on the weeds is essential. Apply only in the Autumn and Winter (mid-April to end of August). For optimum control, apply only when the weeds are actively growing and before the weeds are shielded by the crop.</p> <p>CAUTION: Lucerne and Clover: This product will cause slight leaf burning of undersown lucerne and Clover. DO NOT apply to Persian Clover or Berseem Clover. Linseed: DO NOT spray linseed by aerial application. Lotus: Some initial scorching of leaves may occur in some cases. Medics: Application to some cultivars of Medics may result in unacceptable crop damage in some situations. # Effective control of Paterson's Curse and Fumitories will only be obtained under good growing conditions. ## Where Wireweed is growing under stress, it has been found to be less susceptible.</p>
		Up to the 6 leaf stage or when plants are no more than 50 mm in diameter		2.0 L/ha	
	Mexican Poppy (Argemone ochroleuca), Three-cornered Jack (Doublegee, Spiny Emex), (Emex australis)	Up to the 4 leaf stage or when plants are no			

Situation & Crop	Weeds Controlled	Weed Stage	State	Rate	Critical Comments
As above	Turnip Weed (Giant Mustard), (Rapistrum rugosum)	more than 50 mm in diameter	As above	As above	As above
	Wild Radish (Raphanus raphanistrum), Wild Mustard (Sisymbrium spp.), Wild Turnip (Brassica tournefortii), Wireweed## (Polygonum aviculare), Paterson's Curse# (Salvation Jane) (Echium plantagineum)	As above			
	Fumitories# (Fumaria spp.)		QLD, NSW, VIC, SA, WA, ACT only		
	Pheasant's Eye (Adonis microcarpa) Birds Eye (Carrot Weed) (Cotula australis)	Up to the 4 true leaf stage	SA only	700 mL/ha	DO NOT use where undersown medics occur.
		Greater than the 4 true leaf stage.		1.4 L/ha	
	Fireweed (Senecio spp.)	Seedling	QLD, NSW, ACT only	1.4 L/ha	Apply during the Autumn-Winter period when weed are young and actively growing.
Early flowering stage		2.8 L/ha			
Wheat, Barley, Cereal Rye, Oats and Triticale: including those undersown with Clover, Lucerne or Medics; Clover Seed Crops	Amsinckia (Amsinckia spp.), Capeweed (Arctotheca calendula), Chamomile (Matricaria matricarioides), Charlock (Sinapis arvensis), Corn Gromwell (Ironweed, Sheepweed), (Buglossoides arvensis) Fat Hen	Up to the 5 leaf stage or when weed plants are no more than 50 mm in diameter	WA only	750 mL/ha plus 440 mL of Amicide* 625/ha or plus 700 mL of MCPA 500/ha	CROP STAGE: CEREALS: 3 leaf to fully tillered. (Zadoks Scale Z13-30). CLOVER: Apply after clover seedlings have at least three trifoliate leaves. APPLICATION: Refer to General Instructions. Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application. Apply only in the Autumn and Winter (mid-April to end of August). Apply only when the weeds are actively growing and before the weeds are shielded by the crop.
			VIC, SA only	1.4 L /ha plus	

Situation & Crop	Weeds Controlled	Weed Stage	State	Rate	Critical Comments
As above	(Chenopodium album), Field Madder (Sherardia arvensis), Hexham Scent (King Island Melilot) (Melilotus indicus), Horned Poppy (Glaucium flavum), Indian Hedge Mustard (Sisymbrium orientale), Lesser Swinecress (Coronopus didymus), Mexican Poppy (Argemone ochroleuca), Mintweed (Salvia reflexa), Purple Calandrinia (Calandrinia meziesii) Paterson's Curse (Salvation Jane) (Echium plantagineum) Rough Poppy (Papaver hybridum), Saffron Thistle (Carthamus lanatus), Shepherd's Purse (Capsella bursa-pastoris), Slender Thistle (Carduus tenuiflorus), Turnip Weed (Giant Mustard) (Rapistrum rugosum), Wild Radish (Raphanus raphanistrum)	As above	As above	440 mL of Amicide* 625/ha or plus 700 mL of MCPA 500/ha As above	CAUTION: Some crop damage and/or yield loss may occur in Olympic and Shortim wheat cultivars. DO NOT apply to Persian clover or Berseem clover. A slight burning of clover may occur. DO NOT use 2,4-D or MCPA tank mix if wheat, oats, barley or triticale are undersown with lucerne, clover or Medics unless some damage is acceptable. (continued on next page) This product may be tank mixed with Hoegrass*/Nugrass*.
	Common Peppergrass (Lepidium africanum), Tree Hogweed (Polygonum patulum)		VIC only		
	Black Bindweed (Climbing Buckwheat) (Fallopia convolvulus)	Up to the 8 leaf stage	VIC, SA only		
Wheat, Barley, Oats and Triticale (alone or undersown with Lucerne and	Common Sowthistle (Sonchus oleraceus)				APPLICATION: Refer to General Instructions. Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application. Good coverage of the spray on the weed is essential. Apply

Situation & Crop	Weeds Controlled	Weed Stage	State	Rate	Critical Comments
Clover)					only in the Autumn and Winter (mid-April to end of August). For optimal control, apply when the weeds are actively growing and before the weeds are shielded by the crop. CAUTION: Lucerne and Clover: This product will cause slight leaf burning of undersown Lucerne and Clover. DO NOT apply to Persian Clover and Berseem Clover. (continued below)
		Up to 4 leaves	QLD, Northern NSW	1.4 L/ha	Strictly for use on small weeds early in crop before shading out. CROP STAGE Wheat, Barley, Oats and Triticale: 3-leaf to mid-tillering (Z13-Z23). Use a minimum spray volume of 70L/ha. If Group B resistance is suspected, or weeds have grown through a previous application of chlorsulfuron, apply 2.1L/ha.
		4-8 leaves		2.1 L/ha	CROP STAGE Wheat, Barley, Oats and Triticale: 3-leaf to full-tillering (Z13-Z30). This product will cause slight leaf burning of undersown Lucerne and Clover.
Wheat, Barley, Oats and Triticale	Three-cornered Jack (Doublegee, Spiny Emex) (Emex australis) Wireweed (Polygonum aviculare) Variegated Thistle (Silybum marianum)	Up to the 4 leaf stage or when plants are no more than 50 mm in diameter	QLD, NSW, TAS, SA, ACT only	1.4 L /ha plus 440 mL of Amicide* 625/ha or plus 700 mL of MCPA 500/ha	CROP STAGE: 5 leaf to fully tillered (Zadoks scale Z15-30); Apply under good growing conditions. BOOM SPRAYING: Use a minimum of 220L spray per hectare on Black Bindweed when crop is greater than 300mm high. CAUTION: Some crop damage and/or yield loss may occur in Olympic and Shortim wheat cultivars. DO NOT mix with 2,4-D or MCPA tank mix if wheat, oats, barley or triticale are undersown with lucerne, clover or medics unless some damage is acceptable. DO NOT apply to medics (WA only).
	Fumitories (Fumaria spp.)		QLD, NSW, ACT, VIC only		
	Wild Mustard (Sisymbrium spp.), Wild Turnip (Brassica tournefortii)		QLD, VIC only		This product may be tank mixed with Hoegrass*/Nugrass or control of Annual Ryegrass and Wild Oats in Wheat and Barley only.

Situation & Crop	Weeds Controlled	Weed Stage	State	Rate	Critical Comments
	Cleavers	2-4 stems + 1-3 whorls of leaves /stem	QLD, VIC, NSW, ACT, TAS only		DO NOT use 2,4-D or MCPA tank mix if using a tank with diclofop-methyl. APPLICATION: Refer to General Instructions. Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application.
	Three-cornered Jack (Doublegee, Spiny Emex) (Emex australis) Wireweed (Polygonium)	4 to 5 leaf stage	WA only	2.1 L/ha plus 550 mL of 500 g/L 2,4-D amine or plus 700 mL of 500 g/L MCPA amine	
			QLD, VIC, NSW, ACT, SA, TAS only		
	Fumitories (Fumaria spp.)		QLD, NSW, ACT, VIC, SA only		
	Black Bindweed (Fallopia convolvulus) (Climbing Buckwheat), Chamomile (Matricaria matricarioides), Fat Hen (Chemopodium album), Field Madder (Sherardia arvensis), Lesser Swinecress (Coronopus didymus) Mountain Sorrel Oxalis acetosella) Purple calandrinia (Calandrinia menziesii) Paterson's Curse (Echium plantagineum) (Salvation Jane) Shepherd's Purse (Capsella bursa-pastoris) Three-horned bedstraw (Gallium trichomutum) Variegated Thistle (Silybum marianum)	5 to 8 leaf stage	All states		
Fallow & Non-Crop	Cowvine or Peachvine (Ipomea lonchophylla)	Up to 6 leaves	QLD, NSW, ACT only	1.4 L/ha	<u>For all fallow and non-crop situations</u> Complete coverage is essential, plants which are shaded by stubble or other weeds will NOT be controlled. Weeds that are drought stressed and sprayed under hot conditions may NOT be
		Up to 8 leaves		2.1 L/ha	

Situation & Crop	Weeds Controlled	Weed Stage	State	Rate	Critical Comments
	Climbing Buckwheat (Fallopia convolvulus)	Cotyledon to mature, vining plants		1.5 L/ha alone or 1 – 1.5 L/ha plus glyphosate	adequately controlled. The addition of glyphosate will provide broader spectrum weed control when needed. Refer to the compatibility section for appropriate glyphosate products. The rate of glyphosate used should be robust for the conditions and sufficient to control the hardest-to-kill weed in the situation. For general fallow weed control application volumes above 50L/ha are recommended.
	Volunteer conventional cotton (Gossypium hirsutum)	Cotyledon to 6 leaves	QLD, NSW, WA, NT only	1.5 L/ha alone or 1 – 1.5 L/ha plus glyphosate	For Volunteer Cotton Control, including Roundup Ready* varieties, a minimum spray volume of 80 L/ha should be used. Wellfarm Bromoxynil 200 EC Herbicide alone or with Glyphosate will NOT provide satisfactory control of ratoon or stub cotton.
	Volunteer RoundUp Ready* cotton (Gossypium hirsutum)	Cotyledon to 6 leaves		1.5 L/ha alone or plus glyphosate	
Pastures	Adonis (Pheasants Eye), Birds Eye (Carrot Weed)	Up to 4 leaves	SA only	700 mL/ha	Apply in May-June. Medics may be severely scorched at the low rate and killed at the high rate. DO NOT apply to polymorpha medics. Avoid application to clover-based pastures in temperatures over 20°C or when temperatures above 20°C may follow for some days after application, as seedling mortality may occur and established plants may be damaged.
		More than 4 leaves		1.4L/ha	
	Doublegee, Wild Radish	Up to 4 leaves	WA only	1.5L/ha	
	Matricaria	Up to 3 leaves		750 mL/ha	
	Fireweed	Seedling	NSW, QLD, ACT only	1.4L/ha	Apply with low volume boom spray during Autumn/Winter, when weeds are young and actively growing.
		Early flowering		2.8L/ha	
Pastures, Roadsides, Rights-of-way	African Daisy (Senecio pterophorus)	Up to and including the flowering stage	SA only	300 mL per 100 L water	The plants, including the stems, must be thoroughly wet to the point of run-off. Apply when the plants are actively growing. A non-ionic wetter should be added at 112 g non-ionic surfactant per 100 litres of spray (or at labelled rate). A follow-up treatment may be necessary for large, well-established plants.
	Boneseed/Bitou Bush (Chrysanthemoides monilifera)	Seedling only	VIC, TAS only	160 mL per 100 L water	Ensure an overall spray coverage on weed seedlings.

Situation & Crop	Weeds Controlled	State	Rate			Critical Comments
Turf Established couch, (including QLD Blue), Buffalo, Bent, Paspalum, Kikuyu, Fescue and Rye lawns	Bindy-eye (Calotis hispidula), Jo-jo (Onehunga) (Soliva pterosperma), Cudweed (Gnaphalium spp.)	All States	6L/500 L water per ha	High Volume 300 mL per 100 L water	Knapsack 60 mL/ 100m ² per 15L	Apply evenly over the area to be treated. A slight browning of grasses, especially Paspalum, may occur in hot conditions.

* Other registered trademarks

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

WITHHOLDING PERIODS:

GRAZING	DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 8 WEEKS AFTER APPLICATION.
HARVEST	NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

Before opening, carefully read Directions for Use, Precautionary and Protection Statements, Storage and Disposal, Safety Directions and First Aid instructions. Use only at recommended growth stages of crops and weeds. DO NOT use on crops other than recommended.

MIXING INFORMATION

To ensure even mixing, half fill the spray tank with clean water and add the required amount of this product. Agitate thoroughly, then add the remainder of the water. Agitate again before spraying commences.

BOOM SPRAYER

Apply 50 to 200 litres of water per hectare (WA: 50 to 100L/ha). Calibrate the machine before use to ensure that the correct application rate is obtained. Check height of boom above the crop to ensure that the spray is evenly distributed. A minimum of 160 litres and preferably 200 litres of water per hectare is recommended where the weed infestation is heavy or the crop cover is dense. DO NOT use rain-drop or air-induction nozzles as coverage of weeds may NOT be adequate.

AERIAL APPLICATION

Apply a minimum of 22 litres of water per hectare. Markers should always move upwind before aircraft passes overhead. Application by Micronair equipment may give variable results.

EQUIPMENT MAINTENANCE

After use wash equipment thoroughly and rinse several times prior to re-use. Wash sprayer thoroughly with clean water after use. If using tank mixtures with 2,4-D Amine or MCPA 250 apply with equipment that will NOT be used in 2,4-D and MCPA susceptible crops, particularly cotton and vines.

COMPATIBILITY

This Product May Be Mixed In The Spray Tank With One Of The Following Herbicides And Insecticides In Wheat And Barley Only:

Metsulfuron-Methyl, Chlorsulfuron, 2,4-D Amine, MCPA 250, Diflufenican, Dicamba, Chlorpyrifos and Omethoate, Halosulfuron-Methyl, Glyphosate products, including Wellfarm products.

As formulations of other manufacturer's products are beyond the control of Wellfarm Pty Ltd, all mixtures should be tested prior to mixing commercial quantities.

RESISTANT WEEDS WARNING

GROUP	C	HERBICIDE
--------------	----------	------------------

Wellfarm Bromoxynil 200 EC Herbicide is a member of the nitriles group of herbicides. Wellfarm Bromoxynil 200 EC Herbicide has the inhibitors of photosynthesis at photosystem II mode of action. For weed resistance management, Wellfarm Bromoxynil 200 EC Herbicide is a Group C herbicide. Some naturally occurring weed biotypes resistant to Wellfarm Bromoxynil 200 EC Herbicide and other Group C herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if Wellfarm Bromoxynil 200 EC Herbicide and other Group C herbicides are used repeatedly. These resistant weeds will NOT be controlled by Wellfarm Bromoxynil 200 EC Herbicide or any other Group C herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Wellfarm Pty Ltd accepts NO liability for any losses that may result from the failure of Wellfarm Bromoxynil 200 EC Herbicide to control resistant weeds.

Wellfarm Bromoxynil 200 EC Herbicide may be subject to specific resistance management strategies. For further information consult your local farm chemical supplier, consultant, local Department of Agriculture or Primary Industries.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DRIFT WARNING: DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

NOTE: Maximum daily temperature at or above 20°C at application, or on days following treatment may result in seedling mortality and established plants may be damaged.

PROTECTION OF LIVESTOCK

LOW HAZARD TO BEES.

May be applied at any time as recommended in the Directions for Use Table.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DANGEROUS TO FISH.

DO NOT contaminate dams, streams, drains, rivers or waterways with the chemicals or used containers.

STORAGE AND DISPOSAL

KEEP OUT OF REACH OF CHILDREN

For Only (5L, 20L, 200L)

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty containers for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty containers 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory Government Regulations. DO NOT burn empty containers or product.

For Refillable Containers (1000L)

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Product is poisonous if inhaled or swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing spray, wear elbow-length PVC gloves and face shield. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield.

FIRST AID

If poisoning occurs contact a Doctor or phone the Australian Poisons Information Centre 13 11 26. If swallowed, DO NOT induce vomiting. Give a glass of water.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS) that can be obtained from the supplier.

CONDITIONS OF SALE

The use of this product is beyond the control of the manufacturer. No warranty expressed or implied is given by Wellfarm Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Wellfarm Pty Ltd accepts NO responsibility for any consequences whatsoever resulting from the use of this product.

MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY

Wellfarm Pty Ltd has no control over storage, handling and manner of use of this product. Where the material is not stored, handled or used correctly and in accordance with the directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Wellfarm Pty Ltd accepts NO liability for any losses or damages arising from incorrectly storage, handling or use.

IN AN EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE

APVMA Approval No 70239/62804

Wellfarm Pty Ltd
ACN 158 518 551
22 Calypso Crescent
Point Cook Vic 3030
www.wellfarm.com.au