

CAUTION KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Epoxiconazole 125 SC

Fungicide

ACTIVE CONSTITUENT: 125g/L EPOXICONAZOLE

GROUP 3 FUNGICIDE

For the control of leaf rust, stripe rust and septoria nodorum blotch of wheat and leaf rust of barley, as specified in the Directions For Use Table

**IMPORTANT: READ THE ATTACHED LEAFLET
BEFORE OPENING OR USING THIS PRODUCT**

CONTENTS

10L

STORAGE AND DISPOSAL KEEP OUT OF REACH OF CHILDREN

Store in the closed, original container in a cool, well ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty containers for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty containers 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory Government Regulations. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

Harmful if inhaled or swallowed. Will irritate the eyes, nose and throat and skin. Avoid contact with eyes and skin. DO NOT inhale vapours. If product on skin, immediately wash area with soap and water. Wash hands after use. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and half-piece respirator. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and a washable hat. After each day's use, wash gloves, respirator and if rubber, wash with warm water. Wash contaminated clothing.

FIRST AID

If poisoning occurs contact a Doctor or phone the Australian Poisons Information Centre 13 11 26.



Wellfarm Pty Ltd
ACN 158 518 551
ABN 32 158 518 551

22 Calypso Crescent
Point Cook Vic 3030
www.wellfarm.com.au

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS) that can be obtained from the supplier.

CONDITIONS OF SALE

The use of this product is beyond the control of the manufacturer. No warranty expressed or implied is given by Wellfarm Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Wellfarm Pty Ltd accepts NO responsibility for any consequences whatsoever resulting from the use of this product.

MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY

Wellfarm Pty Ltd has no control over storage, handling and manner of use of this product. Where the material is not stored, handled or used correctly and in accordance with the directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Wellfarm Pty Ltd accepts NO liability for any losses or damages arising from incorrectly storage, handling or use.

APVMA Approval No 70300/62954

**IN AN EMERGENCY
DIAL 000 POLICE
OR FIRE BRIGADE**

Batch No

DOM



CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Wellfarm Epoxiconazole 125 SC Fungicide

ACTIVE CONSTITUENT: 125g/L EPOXICONAZOLE

GROUP	3	FUNGICIDE
-------	---	-----------

For the control of leaf rust, stripe rust and septoria
nodorum blotch of wheat and leaf rust of barley, as
specified in the Directions For Use Table

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE
OPENING OR USING THIS PRODUCT**

**CONTENTS: 10L
(5L, 20L)**

APVMA Approval No 70300/62954

Wellfarm Pty Ltd
ACN 158 518 551
22 Calypso Crescent
Point Cook Vic 3030
www.wellfarm.com.au



DIRECTIONS FOR USE

CROP	DISEASE	RATE PER HA	CRITICAL COMMENTS
Wheat	Leaf rust (Puccinia recondita)	500 mL plus 200mL non-ionic surfactant per 100L spray-mix	Apply when conditions favour disease development and preferably prior to development of high levels of disease in the crop. Aim to apply between stem elongation and ear emergence complete (ZGS 32-59). DO NOT apply later than ZGS 59. Repeat spraying may be required for stripe rust. Regularly monitor the crop from 3-4 weeks after spraying for signs of re-infection and treat accordingly.
	Stripe rust (Puccinia striiformis)		
	Septoria nodorum blotch (Septoria nodorum)		
Barley	Leaf rust (Puccinia hordei)	250 to 500 mL plus 200mL non-ionic surfactant per 100L spray-mix	Apply when conditions favour disease development and preferably prior to development of high levels of disease in the crop. Aim to apply between stem elongation and ear emergence complete (ZGS 32-59). DO NOT apply later than ZGS 59. Apply the higher rate when disease is present on the top leaf or conditions are favourable for disease development at these stages of crop development.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

DO NOT HARVEST FOR 6 WEEKS AFTER APPLICATION.

DO NOT GRAZE OR CUT FOR STOCKFOOD FOR 6 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

Wellfarm Epoxiconazole 125 SC Fungicide is a broad-spectrum contact and systemic fungicide. It is taken up via the stem and foliage and is translocated upwards and outwards to provide protection to new growth. Thorough coverage of the crop is necessary for best results.

Check the crop regularly for any symptoms of disease infection. Apply as soon as disease is present on the flag leaf or flag-2. Application beyond flag leaf emergence (ZGS 39) may NOT provide any yield benefit, particularly under dry seasonal conditions.

MIXING

Wellfarm Epoxiconazole 125 SC Fungicide is a suspension concentrate formulation. Add the product to the half-filled spray tank while agitating. Continue to agitate while topping up the tank and during spraying.

APPLICATION

Ground Application: Apply in a water volume of between 50 and 100L/ha, using flat fan nozzles operating at around 50 cm above the top of the crop. Use the higher water volume in crops with heavier canopies.

Aerial Application: Apply with suitable aircraft set up and operated to apply fungicides to cereal crops in a minimum water volume of 20L/ha.

COMPATIBILITY

For further information consult your farm chemical supplier, consultant, local Department of Agriculture or Primary Industries.

FUNGICIDE RESISTANCE WARNING

GROUP	3	FUNGICIDE
--------------	----------	------------------

Wellfarm Epoxiconazole 125 SC Fungicide is a member of the DMI group of fungicides. For fungicide resistance management, Wellfarm Epoxiconazole 125 SC Fungicide is a Group 3 fungicide.

Some naturally occurring individual fungi resistant to Wellfarm Epoxiconazole 125 SC Fungicide and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if Wellfarm Epoxiconazole 125 SC Fungicide and other Group 3 fungicides are used repeatedly. These resistant fungi will NOT be controlled by Wellfarm Epoxiconazole 125 SC Fungicide or any other Group 3 fungicides. Thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant individuals is difficult to detect prior to use, Wellfarm Pty Ltd accepts NO liability for any losses that may result from the failure of Wellfarm Epoxiconazole 125 SC Fungicide to control resistant fungi.

Wellfarm Epoxiconazole 125 SC Fungicide may be subject to specific resistance management strategies. For further information consult your local farm chemical supplier, consultant, local Department of Agriculture or Primary Industries.

RE-ENTRY PERIOD

DO NOT allow entry into treated areas until the spray has dried.

CAUTION EXPORT OF TREATED CEREALS

Growers should note that Maximum Residue Limits (MRLs) or import tolerances DO NOT exist in all markets for cereals treated with Wellfarm Epoxiconazole 125 SC Fungicide. Additionally, some export markets have established MRLs different to those in Australia. If you are growing cereals for export, please check with your local supplier or your grain exporter for the latest information on MRLs and import tolerances BEFORE using Wellfarm Epoxiconazole 125 SC Fungicide.

LIVESTOCK DESTINED FOR EXPORT MARKETS

The grazing withholding period only applies to stock slaughtered for the domestic market. Some export markets apply different standards. To meet these standards, ensure that in addition to complying with the grazing withholding period, the Export Slaughter Interval is observed before stock are sold or slaughtered.

EXPORT SLAUGHTER INTERVAL (ESI)

AFTER OBSERVING THE PERIOD FOR GRAZING OR CUTTING FOR STOCK FOOD, LIVESTOCK THAT HAVE BEEN GRAZED ON OR FED TREATED CROPS SHOULD BE PLACED ON CLEAN FEED FOR 3 DAYS PRIOR TO SLAUGHTER.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

HIGHLY TOXIC TO AQUATIC ORGANISMS.

DO NOT apply under weather conditions or from spraying equipment which could be expected to cause spray drift onto adjacent crops, croplands, livestock, pastures, wetlands, waterbodies or watercourses.

DO NOT apply Wellfarm Epoxiconazole 125 SC Fungicide aerially if waterbodies, watercourses or wetlands are within 30 metres downwind of the application area.

DO NOT contaminate dams, streams, drains, rivers or waterways with the chemicals or used containers.

STORAGE AND DISPOSAL

KEEP OUT OF REACH OF CHILDREN

Store in the closed, original container in a cool, well ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty containers for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty containers 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots,

in compliance with relevant Local, State or Territory Government Regulations. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

Harmful if inhaled or swallowed. Will irritate the eyes, nose and throat and skin. Avoid contact with eyes and skin. DO NOT inhale vapours. If product on skin, immediately wash area with soap and water. Wash hands after use. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and half-piece respirator. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and a washable hat. After each day's use, wash gloves, respirator and if rubber, wash with warm water. Wash contaminated clothing.

FIRST AID

If poisoning occurs contact a Doctor or phone the Australian Poisons Information Centre 13 11 26.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS) that can be obtained from the supplier.

CONDITIONS OF SALE

The use of this product is beyond the control of the manufacturer. No warranty expressed or implied is given by Wellfarm Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Wellfarm Pty Ltd accepts NO responsibility for any consequences whatsoever resulting from the use of this product.

MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY

Wellfarm Pty Ltd has no control over storage, handling and manner of use of this product. Where the material is not stored, handled or used correctly and in accordance with the directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Wellfarm Pty Ltd accepts NO liability for any losses or damages arising from incorrectly storage, handling or use.

IN AN EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE

APVMA Approval No 70300/62954

Wellfarm Pty Ltd
ACN 158 518 551
22 Calypso Crescent
Point Cook Vic 3030
www.wellfarm.com.au