

POISON KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Diflufenican 25 + Bromoxynil 250 EC

Herbicide

**ACTIVE CONSTITUENT: 250g/L BROMOXYNIL
PRESENT AS THE OCTANOATE 25g/L DIFLUFENICAN
SOLVENT: 416g/L HYDROCARBON LIQUID
150g/L N-METHYL-2-PYRROLIDONE**

GROUP CF HERBICIDE

For the control of certain broadleaf weeds in winter cereals and pastures as specified in the Directions For Use Table

**IMPORTANT: READ THE ATTACHED LEAFLET
BEFORE OPENING OR USING THIS PRODUCT**

CONTENTS

5L

STORAGE AND DISPOSAL KEEP OUT OF REACH OF CHILDREN

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight.

For Only (5L, 10L, 20L, 200L)

Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty containers for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty containers 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory Government Regulations. DO NOT burn empty containers or product.

For Envirodram - Micro Matic Valve (110L Only)

DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodram with water or any other foreign matter. After each use of the product please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodram have been used, please return the empty Envirodram to the point of purchase.

For Refillable Containers (1000L Only)

The container must be vented before discharging contents. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Product is harmful if inhaled or swallowed. Will irritate the eyes, nose, throat and skin. Avoid inhaling spray mist. When preparing spray wear elbow-length PVC gloves and face-shield. If product in eyes, wash it out immediately with water. After use and before eating, drinking, smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face-shield and contaminated clothing.

FIRST AID

If poisoning occurs contact a Doctor or phone the Australian Poisons Information Centre 13 11 26. If swallowed, DO NOT induce

vomiting. Give a glass of water. If in eyes, wash out immediately with water.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS) that can be obtained from the supplier.

CONDITIONS OF SALE

The use of this product is beyond the control of the manufacturer. No warranty expressed or implied is given by Wellfarm Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Wellfarm Pty Ltd accepts NO responsibility for any consequences whatsoever resulting from the use of this product.

MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY

Wellfarm Pty Ltd has no control over storage, handling and manner of use of this product. Where the material is not stored, handled or used correctly and in accordance with the directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Wellfarm Pty Ltd accepts NO liability for any losses or damages arising from incorrectly storage, handling or use.

APVMA Approval No 70423/63430

**NOT SUBJECT TO THE ADG CODE WHEN TRANSPORTED BY ROAD
OR RAIL IN AUSTRALIA, BUT CLASSED AS DANGEROUS
BY IATA AND IMDG WHEN CARRIED BY AIR OR SEA TRANSPORT**

UN NO. 3082

**ENVIRONMENTALLY HAZARDOUS
SUBSTANCES, LIQUID, N.O.S.**

**IN AN EMERGENCY
DIAL 000 POLICE OR FIRE BRIGADE**

PG III

HAZCHEM 3Z



Wellfarm Pty Ltd
ACN 158 518 551
ABN 32 158 518 551

22 Calypso Crescent
Point Cook Vic 3030
www.wellfarm.com.au



Batch No

DOM

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide

ACTIVE CONSTITUENT:
250g/L BROMOXYNIL PRESENT AS THE OCTANOATE
25g/L DIFLUFENICAN
SOLVENT:
416g/L HYDROCARBON LIQUID
150g/L N-METHYL-2-PYRROLIDONE

GROUP	CF	HERBICIDE
-------	----	-----------

For the control of certain broadleaf weeds in winter cereals
and pastures as specified in the Directions For Use Table

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE
OPENING OR USING THIS PRODUCT**

CONTENTS: 5L
(10L, 20L, 110L, 200L, 1000L)

APVMA Approval No 70423/63430

Wellfarm Pty Ltd
ACN 158 518 551
22 Calypso Crescent
Point Cook Vic 3030
www.wellfarm.com.au



DIRECTIONS FOR USE**RESTRAINTS**

DO NOT apply if crops or weeds are stressed due to dry or excessively moist conditions.

DO NOT apply to crops under stress due to disease or insect damage.

DO NOT apply to frost-affected crops or if frosts are imminent.

DO NOT apply if heavy rain is expected within 4 hours.

DO NOT apply with crop oils (cereals only).

CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE/HA	CRITICAL COMMENTS
Wheat, Barley Triticale, Cereal rye (including cereals undersown with clover and/or lucerne), and these cover crops in vineyards PASTURE Clover and/or lucerne based pasture (newly sown or established including cover crops in vineyards)	Wild Radish (Raphanus raphanistrum)	Up to 2 leaf stage and not more than 60mm in diameter and where weed density is less than 50 plants/m ²	WA only	350 mL	CROP STAGE: Cereals 2 leaf to fully tillered (Zadock's Z12-29). Optimum results are achieved when sprayed at 4-8 weeks post-sowing. WARNING: Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide may cause transient crop yellowing of cereals. (Refer to "Crop Tolerance" section of General Instructions). Clover and Lucerne Application is recommended prior to the 8 th trifoliate leaf stage. Application can be made from the 1 st trifoliate leaf stage in QLD, NSW, ACT and VIC only. In other States applications prior to the 3 leaf stage may result in crop damage if seedlings are under stress and in sandy soils. DO NOT apply to annual medics. WARNING: Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide may affect growth and seed set of some varieties of clover and lucerne. (Refer to "Crop Tolerance" section of the General Instructions). COVER CROPS IN VINEYARDS: When using in vineyard situations, apply during vine dormancy only. Contact with vines must be avoided. Particular care should be taken if applied in late autumn or early spring, when vines may NOT be fully dormant.
	Wild Mustard (Sisymbrium spp.), Wild Radish (Raphanus raphanistrum)	Up to 4 leaf stage and not more than 120mm in diameter	All States	500 mL	
		Up to 6 leaf stage and not more than 150mm in diameter		750 mL	
		Up to 8 leaf stage and not more than 180mm in diameter		1.0L	
	Canola (Rapeseed), Charlock (Sinapis arvensis), Turnip Weed (Rapistum rugosum), Wild Turnip (Brassica tournefortii)	Up to 2 leaf stage and not more than 60mm in diameter		500 mL	
		Up to 4 leaf stage and not more than 120mm in diameter		750 mL	
	Shepherd's Purse (Capsella bursa-pastoris)			1.0L	

	Capeweed (<i>Arctotheca calendula</i>)	Up to 4 leaf stage and not more than 120 mm in diameter		500 mL	<p>WEED STAGE: Apply from early post-emergence.</p> <p>APPLICATION: Apply when weeds are actively growing. Ensure thorough coverage of weeds. Where crop or weed density is high, increase water volume.</p> <p>In most situations the rate specified for each weed size will give satisfactory control. However, under certain conditions such as:</p> <ul style="list-style-type: none"> • high crop and weed density. • late season germinations. • abnormal weed growth (including early flowering): higher rates of product (up to the maximum rate of application specified for that weed) may be required.
		Up to 6 leaf stage and not more than 150mm in diameter		750 mL	
		Up to 8 leaf stage and not more than 180mm in diameter		1.0L	
	Corn Gromwell (<i>Buglossoides arvensis</i>)	Up to 4 leaf stage		500 mL	
		Up to 6 leaf stage		750 mL	
	Climbing Buckwheat (<i>Fallopia convolvulus</i>)	Up to 2 leaf stage		500 mL	
		Up to 4 leaf stage		750 mL	
		Up to 6 leaf stage		1.0L	
					Continued on next page.....

CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE/HA	CRITICAL COMMENTS
Wheat, Barley Triticale, Cereal rye (including cereals undersown with clover and/or lucerne), and these cover crops in vineyards PASTURE Clover and/or lucerne based pasture (newly sown or established including cover crops in vineyards)	Deadnettle (Lamium amplexicaule), Paterson's Curse (Salvation Jane), Rough Poppy (Papaver hybridum)	Up to 2 leaf stage	All States	500 mL	Continued from previous page.....
		Up to 4 leaf stage		750 mL	Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide will NOT effectively control: <ul style="list-style-type: none"> • regrowth of suppressed weeds. • transplanted weeds. • regrowth from rhizomes or roots. • weeds growing under stress from previous herbicide applications. • radish plants beyond rosette stage.
	Amsinckia (Amsinckia spp.)				
	Doublegee (Spiny emex)	Up to 2 leaf stage	QLD, NSW, ACT, VIC, TAS, WA only	500 mL	WILD RADISH: Effective residual activity of this product may be reduced where: <ul style="list-style-type: none"> • rates lower than 1.0L/ha are used. • dry conditions prevail. • poor coverage of the soil surface is achieved. • crop is planed in non-wetting sand. • soils contain a high content of organic matter.
		Up to 4 leaf stage	All States	750 mL	
	Chamomile (Matricaria matricarioides), Common Peppergrass (Lepidium africanum) , Lesser Swinegrass (Coronopus didymus), Purple Calandrina (Mountain Sorrel), Tree Hogweed (Polygonum patulum)	Up to 4 leaf stage		1.1L	VOLUNTEER LUPINS: In some situations, the higher rate of 1.0L/ha may be required to effectively suppress volunteer lupins at the 4 leaf stage. # Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide will suppress Seedling Dock but will NOT suppress regrowth from transplanted roots.
	Fat Hen (Chenopodium album), Field Madder (Sherardia arvensis), Saffron Thistle (Carthamus lanatus), Variegated Thistle (Silybum marianum)			1.0L	

	Ox-tongue (<i>Picris echioides</i>), Wireweed (<i>Polygonum aviculare</i>)	Up to 2 leaf stage			
	Fireweed (<i>Senecio</i>)	Up to 4 leaf stage	QLD, NSW, ACT, VIC, SA, WA, NT only	500 mL	
	Common Cotula (Bird's eye.) Pheasants Eye (<i>Adonis</i>)	Up to 4 leaf stage	SA only	560 mL	
		Greater than 4 leaf stage		1.1L	
Wheat, Barley, Triticale, Cereal Rye	Fumitory (<i>Fumaria</i>)	2-6 leaf stage	WA only	350 + 200 mL/ha terbutryn (500 g/L)	

CROP	WEEDS CONTROLLED	WEED STAGE	STATE	RATE/HA	CRITICAL COMMENTS
	SUPPRESSION OF THE FOLLOWING WEEDS				
Wheat, Barley Triticale, Cereal rye (including cereals undersown with clover and/or lucerne), and these cover crops in vineyards PASTURE Clover and/or lucerne based pasture (newly sown or established including cover crops in vineyards)	Dense-flower Fumitory (Fumaria densiflora)	Up to the 2 leaf stage	All States	750 mL	Refer to Critical Comments from previous page.....
	Chickweed (Stellaria media), Common Sowthistle (Milk thistle), Dock (Rumex), Hexham Scent (King Island Melilot), Prickly Lettuce (Lactuca serriola), Scarlet Pimpernel (Anagallis arvensis), Skeleton Weed (Chondrilla juncea) Sorrel (Rumex acetosella), Speedwell (Veronica), Three-horned Bedstraw (Gallium tricornutum) , Toad Rush (Juncus bufonius)	Up to the 4 leaf stage		1.0L	
	Volunteer Lupins (Lupins angustifolius)			500 mL- 1.0L	
	Crassula (Stonecrop) (Crassula spp.)			Up to 5 Leaf stage	
	Long Storksbill (Erodium botrys)	Up to 4 leaf stage			
	Volunteer Field Peas (Pisum sativum)	Up to 5 node stage		750 mL	
	Ward's Weed (Carrichtera annua)	Up to 5 leaf stage		1.0L	

	Vetch (<i>Vicia sativa</i>)	Up to 2 leaf stage			
	Mouse-eared Chickweed (<i>Cerastium glomeratum</i>)		NSW & ACT only		
	Mexican Poppy (<i>Argemone ochroleuca</i>)		QLD only		
	Mintweed (<i>Salvia reflexa</i>), Spoon Cudweed (<i>Stuartina muelleri</i>)	Up to 4 leaf stage	NSW & ACT only		
	New Zealand Spinach (<i>Tetragonia tetragonoides</i>)	Up to 2 leaf stage	QLD only	750 mL	
	Cleavers (<i>Galium aparine</i>)	Up to 1 whorl stage	SA only	1.0L	
	Ball Mustard (<i>Neslia paniculata</i>)	Up to 4 leaf stage			
	Horehound (<i>Marubium vulgare</i>)	Pre-emergence			
	Marshmallow (<i>Malva parviflora</i>)	Up to 2 leaf stage			

CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE/HA	CRITICAL COMMENTS
Wheat, Barley, Triticale, Cereal rye	Wild Radish (Raphanus raphanistrum)	Up to 4 leaf stage and not more than 120mm in diameter	WA only	350 mL plus 200 mL MCPA LVE (500g/L)	Refer also to all Critical Comments for cereals above. DO NOT use this tank-mix if cereals are undersown with lucerne or annual medics.
		Up to 6 leaf stage and not more than 150mm in diameter	All States	500 mL plus 200 mL MCPA LVE (500g/L)	DO NOT use this tank-mix in vineyards. Crop Stage
		Up to 8 leaf stage and not more than 180mm in diameter.		500 mL plus 400 mL MCPA LVE (500g/L)	Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide 350 mL + MCPA LVE 200 mL: Apply from 3 leaf to fully tillered (Zadok's Z13 to Z30). Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide 500 mL + MCPA LVE 200 mL: Apply from 3 leaf to fully tillered (Zadok's Z13 to Z30). Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide 500 mL + MCPA LVE 400 mL: Apply from 5 leaf stage to fully tillered (Zadok's Z15 to Z30). Optimum results are achieved when sprayed at 4-8 weeks post sowing. WARNING: Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide may cause transient crop yellowing of cereals. (Refer to "Crop Tolerance" section of General Instructions). Observe instructions also on MCPA LVE product label.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

HARVEST		
	CEREALS, GRAPES	NOT REQUIRED WHEN USED AS DIRECTED.
GRAZING		
	PASTURE, CEREALS	DO NOT GRAZE OR CUT FOR STOCK FOOD WITHIN 14 DAYS AFTER APPLICATION.

GENERAL INSTRUCTIONS

- Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide is a post-emergence contact herbicide, which may provide residual control of wild radish up to 4 weeks after application.
- Apply Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide immediately after mixing. DO NOT allow to stand in the spray tank over night.
- Optimum results will be obtained if good soil moisture exists at an dafter application and weeds are NOT stressed.
- Some pre-emergence herbicides, such as atrazine, can cause stress to certain crops resulting in an increase in crop damage when using this product. Crops which are particularly sensitive are lucerne and subterranean clover.

RESISTANT WEEDS WARNING

GROUP	C F	HERBICIDE
--------------	------------	------------------

Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide is a member of the nitrile and nicotinanilide groups of herbicides. Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide is an inhibitor of photo synthesis at photo system II and carotenoid biosynthesis. For weed resistance management, Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide is both a Group C and a Group F herbicide.

Some naturally occurring weed biotypes resistant to Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide and other Group C and Group F herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide and other Group C and Group F herbicides are used repeatedly. These resistant weeds will NOT be controlled by Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide or any other Group C and Group F herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Wellfarm Pty Ltd accepts NO liability for any losses that may result from the failure of Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide to control resistant weeds.

Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide may be subject to specific resistance management strategies. For further information consult your local farm chemical supplier, consultant, local Department of Agriculture or Primary Industries.

TEMPERATURE WARNING

DO NOT apply Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide if frosts are imminent. Frost causes stress on crops and weeds and could result in increased crop effects and/or decreased weed control. To ensure good results Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide should only be applied once the weeds and crop are NO longer under stress from the frost conditions.

Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application, as increased crop damage may result.

CROP TOLERANCE

Cereals

After application some transient crop yellowing may occur. This usually appears as yellow or white banding on leaves. Provided the crop is NOT under stress from pre-emergent herbicide, root disease, insect damage, frost, dry or excessively moist conditions, the development of the crop and subsequent growth will be unaffected.

Lucerne Warning

The tolerance of Lucerne varieties to Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide can vary with rate of application, soil type, crop health, stage of growth and degree of moisture and temperature stress. Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide may result in transient crop yellowing and suppression of growth with a resultant initial reduction in dry matter. For this reason we recommend application prior to the 8 trifoliate leaf stage. However, under normal growing conditions subsequent growth and seed yield should NOT be affected. Crop damage may be increased if rates higher than 500mL/ha are used and in areas where spray over lapping has occurred.

Under normal growing conditions, the following Lucerne varieties have shown acceptable levels of foliage tolerance to Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide applied at 500mL/ha: Hunter River, Nova and Dekalb 185.

Varieties NOT listed should be tested before using Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide over large areas. Consult your farm chemical supplier, consultant, local Department of Agriculture or Primary Industries, or local Wellfarm Pty Ltd representative for advice on specific varieties.

Subterranean Clover

Warning

The tolerance of subterranean clover varieties to Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide can vary with rate of application, soil type, crop health, stage of growth and degree of moisture and temperature stress. Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide may result in transient crop yellowing and suppression of growth with an initial reduction in dry matter. For this reason we recommend application prior to the 8 trifoliate leaf stage. However, under normal growing conditions subsequent growth and seed yield should NOT be affected. Crop damage may be increased if rates higher than 500 mL/ha are used and in areas where spray overlapping has occurred.

Under normal growing conditions, the following varieties have shown acceptable levels of foliage tolerance to Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide applied at 500 mL/ha:

Daliak, Dalkeith, Denmark, Goulburn, Karridale, Leura, Mt. Barker, Nungarin, Rosedale, Seaton Park, Trikkala and Woogenellup.

The variety Junee has shown increased sensitivity to Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide so care should be taken if this variety is part of the pasture sward.

The effects of Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide on subterranean clover seed yield have been tested on the following varieties. Under normal growing conditions they show acceptable levels of tolerance to Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide applied at 500 mL/ha. However, higher rates may reduce seed yield under conditions of low weed pressure: Denmark, Goulburn, Larissa, Nungarin, Seaton Park, Trikkala and Woogenellup.

Varieties NOT listed should be tested before using Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide over large areas. Consult your farm chemical supplier, consultant, local Department of Agriculture or Primary Industries, or local Wellfarm Pty Ltd representative for advice on specific varieties.

Other Clovers

Warning

The tolerance of clover varieties to Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide can vary with rate of application, soil type, crop health, stage of growth and degree of moisture and temperature stress. Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide may result in transient crop yellowing and suppression of growth with a resultant initial reduction in dry matter. For this reason we recommend application prior to the 8 trifoliate leaf stage. However, under normal growing conditions subsequent growth and seed yield should NOT be affected. Crop damage may be increased if rates higher than 500 mL/ha are used and in areas where spray overlapping has occurred. The effect on seed yield of other clovers has NOT been determined.

The following varieties of clover have shown increased sensitivity to Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide:

Big Bee, Sacromonte (Berseem), Haifa (White), Zulu (Arrowleaf), Kyambro, Lupers and Maral (Persian).

Care should be exercised if these clovers are part of the pasture sward.

Varieties NOT listed should be tested before using Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide over large areas. Consult your farm chemical supplier, consultant, local Department of Agriculture or Primary Industries, or local Wellfarm Pty Ltd representative for advice on specific varieties.

Subsequent Crops

To reduce effect on subsequent susceptible crops (e.g. canola), ensure thorough cultivation of soil prior to the sowing of these crops.

MIXING

To ensure even mixing, half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly then add the remainder of the water. Agitate thoroughly while carrying out spray operations. Reseal part used container immediately after use.

APPLICATION

Boom Sprayer

A minimum of 50L water/ha should be used, however, for optimum results water rates of 70-100 L/ha are recommended. Increase the water volume where weed infestation is heavy or the crop cover is dense. Complete coverage of weeds is essential. Higher water volumes (up to 100 L/ha) will ensure faster activity of the product on the weeds but may increase the symptoms of crop damage.

The following settings are examples which will ensure excellent coverage of exposed weeds:

Water Rate	50 L/ha	75 L/ha	75 L/ha
Nozzle	Hardi No. 10 or equivalent	Hardi No. 12 or equivalent	Hardi No. 14 or equivalent
Speed	10 km/h	10 km/h	12 km/h
Pressure	240 kPa (2.4 bar)	220 kPa (2.2 bar)	210 kPa (2.1 bar)

Controlled Droplet Application (CDA)

Insufficient information is available to recommend the application of this product by CDA.

Warning

The rubber components present in some spraying units may be affected by exposure to the solvents in Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide. To reduce this risk it is recommended that the spray unit be thoroughly washed with a boom cleaner and fresh water after use.

Aircraft

Insufficient information is available to recommend the application of this product by air.

COMPATIBILITY

The following herbicide products are physically compatible with Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide as two-way mixtures in the spray tank, but should only be used for the crops specified, and only when the crop is also specified on the label of the compatible product: (See below for list of compatible insecticides).

Crop	Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide	Compatible Product
Wheat, triticale, cereal rye (including undersown)	Up to 750 mL/ha	Hoegrass® (barley also), Tristar® Advance (barley also), Wildcat® 110 EC (wild oats only, high rate)
Wheat, barley, triticale, cereal rye (including undersown)	All rates	Broadstrike®
Wheat, barley, triticale, cereal rye (not undersown)	Up to 500 mL/ha	Ally®, Glean®, MCPA LVE (500 g/L product) (up to 500 mL/ha only)
	All rates	2,4-D Amine 500, Eclipse®, Cadence® (up to 115 g only), Lontrel®
Wheat only (not undersown)		Topik®
Established lucerne only	Up to 750 mL/ha	Simazine (500 g/L product) (up to 1.25 L/ha only) and Simazine (500 g/L)/Paraquat (200 g/L) mixture
Newly sown and established lucerne and clover only	Up to 750 mL/ha	Targa®, Fusilade®, 2,4-DB Amine (500 g/L)
	Up to 1.0 L/ha	Broadstrike®

* Decis Options®, Eclipse®, Hoegrass®, Jaguar®, Thiodan®, Tristar® and Wildcat® are Registered Trademarks of Bayer.

When mixing Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide with other herbicides, crop yellowing may be enhanced.

When mixing with Hoegrass®, Wildcat® 110 EC or Tristar® Advance some reduction in the efficacy and speed of action of these products may occur.

When mixing with Targa® or Fusilade® some reduction in the efficacy and speed of action of these products and Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide may occur.

In tank-mixtures with Ally® and Glean®, rates of Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide higher than 500 mL/ha may cause significant crop damage. If the crop is stressed, the application of the herbicide tank-mixtures may cause yield reduction. When mixing with Cadence® a temporary wilting may be evident in some crops after application. The mixture of Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide and Simazine should be applied during winter to lucerne which is NOT actively growing. This mixture may result in an increased crop effect but this can be reduced if the lucerne is grazed or cut before spraying. DO NOT mix Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide with Verdict®. Growers should seek advice before spraying recently released cereal varieties. Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide may be mixed in the spray tank with one of the following insecticides according to the directions for the insecticide product: Chlorpyrifos (500 g/L product), Decis Options®, Dimethoate, Dominex®100EC, Fastac® Duo, Le-mat® 290 SL, Talstar® and Thiodan®. Use the recommended rates for Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide and its tank-mix partner as well as the surfactant recommendation of the tank-mix partner. Read the label of the tank-mix partner before mixing and using the tank mixture. If another herbicide is applied as a tank mix, observe the plant back restrictions on that label.

Warning

DO NOT use crop oils with Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide or Wellfarm Diflufenican 25 + Bromoxynil 250 EC Herbicide tank mixtures in cereals. As formulations of other manufacturers' products are beyond the control of Wellfarm Pty Ltd, all mixtures should be tested prior to mixing commercial quantities.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Wash sprayer thoroughly after use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DANGEROUS TO FISH.

DO NOT contaminate dams, streams, drains, rivers or waterways with the chemicals or used containers.

STORAGE AND DISPOSAL

KEEP OUT OF REACH OF CHILDREN

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight.

For Only (5L, 10L, 20L, 200L)

Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty containers for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty containers 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory Government Regulations. DO NOT burn empty containers or product.

For Envirodrum - Micro Matic Valve (110L Only)

DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodrum with water or any other foreign matter. After each use of the product please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase.

For Refillable Containers (1000L Only)

The container must be vented before discharging contents. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Product is harmful if inhaled or swallowed. Will irritate the eyes, nose, throat and skin. Avoid inhaling spray mist. When preparing spray wear elbow-length PVC gloves and face-shield. If product in eyes,

wash it out immediately with water. After use and before eating, drinking, smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face-shield and contaminated clothing.

FIRST AID

If poisoning occurs contact a Doctor or phone the Australian Poisons Information Centre 13 11 26. If swallowed, DO NOT induce vomiting. Give a glass of water. If in eyes, wash out immediately with water.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS) that can be obtained from the supplier.

CONDITIONS OF SALE

The use of this product is beyond the control of the manufacturer. No warranty expressed or implied is given by Wellfarm Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Wellfarm Pty Ltd accepts NO responsibility for any consequences whatsoever resulting from the use of this product.

MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY

Wellfarm Pty Ltd has no control over storage, handling and manner of use of this product. Where the material is not stored, handled or used correctly and in accordance with the directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Wellfarm Pty Ltd accepts NO liability for any losses or damages arising from incorrectly storage, handling or use.

IN AN EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE

NOT SUBJECT TO THE ADG CODE WHEN TRANSPORTED BY ROAD OR RAIL IN AUSTRALIA, BUT CLASSED AS DANGEROUS BY IATA AND IMDG WHEN CARRIED BY AIR OR SEA TRANSPORT	
UN NO. 3082	ENVIRONMENTALLY HAZARDOUS SUBSTANCES, LIQUID, N.O.S.
IN AN EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE	
PG III	HAZCHEM 3Z

APVMA Approval No 70423/63430

Wellfarm Pty Ltd
ACN 158 518 551
22 Calypso Crescent
Point Cook Vic 3030
www.wellfarm.com.au